

# Chronology and distribution of Pleistocene woolly rhinoceros: A review of the archival data from Poland

Kamilla Pawłowska\*, Kajetan Dedła, Katarzyna Płonka

Adam Mickiewicz University, Poznań, Institute of Geology, ul. Krygowskiego 12, 61-680  
Poznań, Poland

\*corresponding author, e-mail: koka@amu.edu.pl

## Abstract

This work provides a review of the state of knowledge of woolly rhinoceroses in Poland. We compile research results from the 19th century to the present day and consider the collected data not only quantitatively, but above all qualitatively. Here we present a list of 215 sites from Poland where remains of the woolly rhinoceros have been found. Studies of woolly rhinos from Poland usually employ small samples. Our compilation of data also reveals that there is currently no basis for drawing conclusions regarding the geographical distribution pattern of the species in Poland. Only a small number of works have focused on matching the places where remains occurred with the geological conditions of the area and their depositional history. Moreover, the results show that the resolution of the radiocarbon dates that are currently published is insufficient to allow conclusions about the chronology of woolly rhinoceroses in Poland to be drawn. No works to date have dealt with any aspect of palaeopopulation research. The woolly rhinoceros is not present in Palaeolithic art in Poland. A summary of our knowledge of this taxon is the starting point for our multi-aspect research into this topic.

**Key words:** History, palaeogeography, taphonomy, Central Europe, WOOLRHINOPOLI Project

## 1. Introduction

Rhinoceroses are mammals belonging to the order Ungulates (Perissodactyla), in which the third toe is the most developed and carries the axis of the limb. The term “rhinoceros” covers a number of living and extinct species, with *Stephanorhinus etruscus*, *Stephanorhinus*

32 *hundsheimensis*, the steppe rhino (*Stephanorhinus hemitoechus*), the forest rhino  
33 (*Stephanorhinus kirchbergensis*), and the woolly rhinoceros (*Coelodonta* species; Fig. 1)  
34 recognized from the fossil record of Pleistocene strata. The fifty-million-year history of this  
35 group of animals is now changing drastically with a population decline in the five living species  
36 of rhinoceroses — the white rhinoceros (*Ceratotherium simum*), the black rhinoceros (*Diceros*  
37 *bicornis*), the Indian rhinoceros (*Rhinoceros unicornis*), the Javan rhinoceros (*R. sondaicus*),  
38 and the Sumatran rhinoceros (*Dicerorhinus sumatrensis*) — which together number less than  
39 30,000 individuals today.

40 Fossil members of rhinoceroses are known from Europe, Asia, North America, and  
41 Africa, and studies of their remains have shed light on intraspecies and interspecies  
42 distributional patterns, morphology, genetics, extinction, diet, and other issues.

43 One recently completed morphological study employed a Bayesian model to reveal the  
44 genus of *Stephanorhinus* as a monophyletic clade (Pandolfi, 2023). A consistent finding in  
45 morphological studies is that it is difficult to distinguish between fossil species using most  
46 skeletal elements, particularly postcranial elements, and this is particularly true in the genus  
47 *Stephanorhinus*. This has to do with the fact that interspecific variability often overlaps and  
48 includes polymorphic characters (Ballatore & Breda, 2019).

49 Genetic investigation has shown that the extinct woolly rhinoceros is most closely  
50 related to the extant Sumatran rhinoceros (Orlando et al., 2003). Phylogenetic inference,  
51 based on a complete mitochondrial genome sequence of fossil *Stephanorhinus* from the  
52 Chondon River valley (Arctic Yakutia, Russia), has confirmed that of the extinct species, woolly  
53 rhinoceros, *Coelodonta* is most closely related to *Stephanorhinus* (Kirillova et al., 2017). A  
54 study of ancient DNA from eastern European sites further suggests that the effective  
55 population size of woolly rhinoceros increased at 29.7 ka BP and then remained stable until  
56 near extinction, ca 14 ka BP (Lord et al., 2020). The extinction of the *Stephanorhinus* species  
57 has also been studied, leading to the conclusion that *Stephanorhinus etruscus* went extinct at  
58 the Early–Middle Pleistocene (in Italy and the Iberian Peninsula) or the late Early Pleistocene  
59 (in Central Europe) (Pandolfi et al., 2017), while *Stephanorhinus hundsheimensis* became  
60 extinct in the late Middle Pleistocene–early Late Pleistocene (in Spain) (García-Fernández et  
61 al., 2023), *Stephanorhinus hemitoechus* at 41 ka BP (in Italy) (Pandolfi et al., 2017), and  
62 *Stephanorhinus kirchbergensis* at about the early Late Pleistocene — that is 120-104 ka BP (in  
63 Europe) or 13 ka BP (in China) (Shpansky, 2017).

64 The diet of fossil rhinoceros can be determined using various, often combined,  
65 methods including morphometry, microwear and mesowear analysis of teeth, isotopic  
66 studies, and analysis of organics in the form of food remains extracted from teeth. The dietary  
67 evolution of the two Middle to Late Pleistocene rhinoceros species *S. hemitoechus* and *S.*  
68 *kirchbergensis* in Central and Northwest Europe was traced by van Asperen & Kahlke (2015)  
69 using mesowear signal and morphometry. They showed a mixed feeder diet for both,  
70 comparable with that of extant mammal species in relatively open habitats. *S. kirchbergensis*  
71 consumed more or less browse in the diet depending on the quality of the habitat, while *S.*  
72 *hemitoechus* shifted from a mixed feeder to consuming more grass when necessary (van  
73 Asperen & Kahlke, 2015).

74 Ballatore (2016) has combined powder X-ray diffraction, carbon isotope geochemistry,  
75 tooth wear analysis, and biometry to perform a palaeoecological investigation of Pleistocene  
76 European Rhinoceroses (*Stephanorhinus*), also concluding that *Stephanorhinus* shows high  
77 specialization in browsing. Further work has also confirmed a mixed feeder diet for the forest  
78 rhino (from the Chondon River) through an analysis of food remains in the fossae of the cheek  
79 teeth, identified as *Larix*, *Vaccinium*, *Betula* spp., *Aulacomnium*, and dicotyledonous herbs and  
80 grasses (Kirillova et al., 2017). In addition, microwear analysis of teeth in this study further  
81 showed that, during the last months of its life, this individual fed predominantly on leaves and  
82 twigs (Kirillova et al., 2017). The study of the *Stephanorhinus kirchbergensis* find from the  
83 Chondon River also shows the validity of undertaking complex studies of rhino remains, which  
84 in this case combined genetic, dietary, and morphological analyses (morphometry, food  
85 remains from teeth, mesowear, and isotopes) to determine the taxonomic position, season,  
86 and age of death, as well as the discussed above diet of this individual.

87 Studies combining mesowear and isotope analysis have also confirmed a generally  
88 browse-dominated diet for *S. kirchbergensis* (Pushkina et al., 2020), as also other work  
89 (Stefaniak et al., 2021). This shows how different lines of evidence in *Stephanorhinus* research  
90 have led to the same consistent conclusions. However, the correlation between  $\delta^{13}\text{C}$  and  
91 mesowear is not straightforward, because of the considerable variation in dietary  
92 specialisation within herbivorous mammals. For example, variation in feeding behavior among  
93 the *S. hemitoechus* and *S. kirchbergensis* species was induced by interspecific competition  
94 along with diversification of habitat within the range of each species' ecological tolerance.  
95 While their dietary flexibility enabled them to survive in a range of environments that were

96 neither strictly steppe nor strictly forest, it does also cause difficulties in establishing the  
97 specific habitats and real dietary traits of the *Stephanorhinus* palaeopopulation (van Asperen  
98 & Kahlke, 2015).

99 The woolly rhinoceros (*C. antiquitatis* Blumenbach 1799) has been demonstrated to be  
100 a specialised grazer on the basis of the high mesowear signal resulting from dental analysis.  
101 This points to abrasion-dominated diets and open grassland habitats (Pushkina et al., 2020).

102 Although the woolly rhinoceros provides the most evidence for the presence of fossil  
103 rhinos in Europe, there is still in many respects no synthesis for this taxon. Here, we will focus  
104 on woolly rhinoceros outlining the current state of knowledge, especially from the perspective  
105 of its spatial and temporal occurrence in Poland, by closely scrutinized the results to date on  
106 woolly rhino remains. It will prove extremely valuable to compile for the first time the scopes  
107 of various types of research into the woolly rhinoceros in Poland, by tracking geographical,  
108 chronological, demographic, genetic, and taphonomic datasets to reconstruct its natural  
109 history, as well as social aspects of its history.

110

## 111 **1.1. Woolly rhinoceroses**

112

### 113 **1.1.1. Emergence and evolution of the genus *Coelodonta***

114 The fossil record of the genus *Coelodonta* goes back to 3.7 Ma BP and was established on the  
115 basis of the remains of *Coelodonta thibetana*, found in Tibet (Zanda Basin) in the Pliocene  
116 strata (Deng et al., 2011). Between 2.5 and 1.0 Ma BP the genus is represented by the remains  
117 of *Coelodonta nihowanensis*, found at many localities in China.

118 *Coelodonta* seems to have arrived in Europe in the early Middle Pleistocene, around  
119 500–400 ka, as shown by finds of *Coelodonta tologoijensis*, a species of rhino with more  
120 evolved features, at localities in Germany (Bad Frankenhausen), Russia, and Mongolia  
121 (Nalaikha). However, some authors have claimed that finds from Germany should be assigned  
122 to *C. antiquitatis praecursor* (Uzunidis, 2022). Finds from the Kuznetsk Basin (Russia) recorded  
123 as *C. cf. tologoijensis* indicate an initial westward expansion of this group of rhinoceroses into  
124 south-west Siberia (Kahlke & Lacombat, 2008).

125 Woolly rhinoceros (*Coelodonta antiquitatis*) comparable to *C. tologoijensis* spread  
126 westward and entered Central Europe and, in several cases, Western Europe, during all of the  
127 subsequent Middle to Late Pleistocene cold stages (around 470–350 ka), as a result of

128 preferable environmental conditions — namely, extended phases of low temperature and  
129 aridity (Kahlke & Lacomat, 2008). By the end of the Pleistocene (until ca. 40 ka; Stuart &  
130 Lister, 2012) the woolly rhinoceros was widely distributed geographically in Eurasia.

131

### 132 **1.1.2. Environment and diet**

133 The natural environment is the complex of physical, chemical, and biotic factors (such as  
134 climate, soil, and living things: i.e. Wolfhagen et al., 2020) that act upon an organism or an  
135 ecological community, which ultimately determine its form and survival. Despite the fact that  
136 animals are embedded in specific environments, they possess some ecological flexibility to  
137 adapt when that environment changes.

138 For the woolly rhinoceros, the natural habitat was the steppe-tundra, a biome that  
139 formerly existed but which has no analogue in today's landscapes as its characteristic  
140 vegetation no longer exists. One of the earliest large species of herbivore to appear there was  
141 the steppe mammoth (*Mammuthus trogontherii*), recognized in Poland at fifteen sites  
142 (Pawłowska, 2015a), including Bełchatów, where cultural traces on a rib, likely left by *Homo*  
143 *heidelbergensis* (indirect evidence given the lack of human remains), constitute the oldest  
144 butchery marks from Poland (Pawłowska et al., 2014; Pawłowska, 2017a) and one of a  
145 relatively few from Eurasia (Pawłowska, 2017a).

146 During the peak of the LGM, the mammoth steppe stretched from the Iberian  
147 Peninsula across Eurasia into Alaska and Canada (to Yukon) and served as the main source of  
148 protein for megafauna, which are a subset of the largest terrestrial species in a community or  
149 an ecosystem. However, environments in the Quaternary varied by the presence of  
150 subenvironments which included refugia, ecological niches that are closely related to the  
151 latitude. Environmental fluctuations during the Pleistocene were driven by climate, by  
152 alternating cycles of cold and warming in relation to glacial and interglacial periods, which in  
153 turn occurred in response to the development and disappearance of ice sheets in Europe  
154 (Hrynowiecka et al., 2022). These dynamic environmental changes affected the spatial range  
155 of fauna in the Pleistocene in Europe, including that of the woolly rhinoceros.

156 The main adaptations of the woolly rhinoceros to the extremely cold, harsh conditions  
157 that prevailed on the mammoth steppe included its long, thick hair, horns that supported  
158 recovering food from the snow cover, and genetic mutations that allowed a type of receptor  
159 in the skin for sensing warm and cold temperatures (Fortelius, 1983; Lord et al., 2020). The

160 woolly rhinoceros fed on green plants, mostly grass with shrubs and woody vegetation in the  
161 winter diet, as part of seasonal changes, as shown by isotopic signatures (Tiunov & Kirillova,  
162 2010). A study of the stomach contents of frozen rhinoceroses revealed that the last meals of  
163 those individuals contained cereal grain (grasses) and sedge (Vereshchagin & Baryshnikov,  
164 1992). It is assumed that the low grass density and lack of suitable habitat prevented it from  
165 crossing the Bering Strait to the American continent (Boeskorov, 2001; Prothero, 1993). As a  
166 result, the remains of the woolly rhinoceros are known only in Europe and Asia.

167

### 168 **1.1.3. Fossil records**

169 Woolly rhinoceros fossils are fairly common and have been discovered in a range from  
170 western Europe to northeastern Siberia. Their absence in specific locations seems to indicate  
171 that they did not reach north–central Siberia or North America.

172 Extinct rhinos from Germany, the Czech Republic, Slovakia, Ukraine, and Belarus,  
173 Lithuania, all of which are neighbors of Poland have been the subject of many studies. Geraads  
174 et al. (2021) provide for the first time a comprehensive database of fossil rhinos of the old  
175 world Neogene and Quaternary, showing that representatives of this group were present in  
176 many localities: at 268 sites in Germany (with references to almost 60 works), at 208 sites in  
177 the Czech Republic (with references to almost 20 works), at 34 sites in Slovakia (with  
178 references to almost 70 works), at 32 sites in Ukraine (with references to almost 20 works), at  
179 2 sites in Belarus (with references to 1 work), and at 1 sites in Lithuania (with references to 2  
180 works). They also showed the distribution pattern of fossil rhinos in Central Europe  
181 corresponding to the woolly mammoth, which is comprehensible given the general ecological  
182 requirements of these herbivores.

183 Despite its considerable spatial distribution, the degree of preservation of remains of  
184 *C. antiquitatis* varies, with skeletal elements without articulation predominating. One  
185 exception is the complete specimen of a woolly rhinoceros individual, along with soft tissues,  
186 which was preserved in the fossil record thanks to favorable ozokerite (earth wax) and rock  
187 salt as fossilization conditions that are abundant in the area of the find in Starunia, Ukraine  
188 (Kubiak, 1994; Kowalski, 2000; Kubiak & Drygant, 2005). Well-preserved remains also have  
189 been found in permafrost in Russia (Belyaev et al., 2023; Boeskorov et al., 2011).

190

191 **2. Material and methods**

192

193 The material for the study consists of the remains of woolly rhinoceroses found at Polish sites.  
194 The list of sites that these remains were recovered from is based on works published since the  
195 nineteenth century. For all archival sites, basic data (mainly the location) have been verified  
196 and in some cases have been corrected against Stefaniak et al. (2023) in order to make them  
197 useful in the synthesis. To this end, a new system for specifying sites is proposed here, as was  
198 done in our earlier synthesis of sites with mammoth remains (Pawłowska, 2015a). This means,  
199 among other things, separating the name and context of site, giving the district in the  
200 subheading of the name of site, and the county and voivodship as separate data. This allows  
201 a standardised way of determining site location. Sites with the woolly rhinoceros remains are  
202 given in the order of the authors who discovered or first described them, as such paper  
203 brought new value to the field. We did not include papers that merely duplicate data.

204

205 **3. Results and discussion**

206

207 Like the woolly mammoth, woolly rhinoceros is part of the *Mammuthus–Coelodonta* faunal  
208 complex, a term that refers to cold-adapted Pleistocene large mammal assemblages with  
209 similar or identical faunistic structures, known for their transregional expansion in Eurasia  
210 (Kahlke, 2014). The Middle Pleistocene and Late Pleistocene faunal complexes include various  
211 animal species, which are associated with the evolution of the fauna during those times. The  
212 faunal complex is, generally speaking, made up of animals associated with both the steppe  
213 and tundra.

214 The woolly rhinoceros is a common species found in Upper Pleistocene assemblages in  
215 Poland, as demonstrated by numerous Pleistocene faunal assemblages (see Appendix 1). It is  
216 therefore surprising that there has been little attention paid to research into the remains of  
217 this species from Polish contexts. To overcome this, a multiproxy study has been conducted  
218 on the rhinoceros remains, along with other taxa, from Krosinko in Wielkopolska, which  
219 included taxonomic, taphonomic, radiocarbon, genetic, and social aspects (Pawłowska, 2022,  
220 2023; Pawłowska et al., 2022). The results of these earlier studies, along with the history of  
221 research on the woolly rhinoceros in Poland, will be presented in terms of geographic,  
222 chronological, demographic, genetic, and taphonomic evidence, as well as artistic

223 representation. This site, along with others (such as Sitkówka) have also contributed to  
224 syntheses already compiled for Central Europe (Puzachenko et al., 2022).

225

### 226 **3.1. Geographic and historic evidences**

227

228 The oldest palaeontological discoveries of woolly rhinoceros sites date back to the nineteen  
229 century (Table 1). Although Stefaniak et al. (2023) states that the first summary paper was  
230 published in 1884 by A. Slósarski, who reported on the skull and mandible of a woolly  
231 rhinoceros found near Kamieńczyk in 1815, in fact the first paper that dealt with the synthesis  
232 of Pleistocene fauna was by Staszic (1815) with the telling title “First treatise on...”. This work  
233 described one of the first rhinoceros finds from Poland, which was a mandible found in 1810  
234 near Kamieńczyk, where the Liwiec River reaches the Bug River (Staszic, 1815) (Fig. 2).

235 The results of these studies from the turn of the century were summarised by Kowalski  
236 (1959), who gave a list of almost a hundred sites (Kowalski, 1959) and provided a distribution  
237 of these sites for Poland. The sites are distributed almost all over the country, mainly along  
238 rivers and their tributaries. This list of Polish sites with woolly rhinoceros remains includes the  
239 results of seminal works of Lubicz-Niezabitowski (1913, 1926, 1929, 1938), a summary of  
240 which has been presented by Pawłowska (2015b), along with findings and evaluations of  
241 specimens held in collections. Later studies of the Pleistocene fauna added more sites with  
242 evidence of the occurrence of woolly rhinoceros to the general list, or made use of data from  
243 these all authors (Kubiak, 1969, 1989; Kubiak & Dziurdzik, 1973; Kuc et al., 2005; Kubiak, 2009;  
244 Kuc et al., 2012; Borsuk-Białynicka, 1973). The monograph of Borsuk-Białynicka provided a  
245 morphometric study of skulls from Poland, Russia, and the Czech Republic.

246 Since 1973, rhinos have not disappeared from the Polish research agenda, and have  
247 been the subject of publications (Karaszewski, 1980; Król, 1998; Kaczmarek, 2004;  
248 Wiszniowska et al., 2005; Wojtal, 2007; Woroncowa-Marcinowska et al., 2013; Pawłowska,  
249 2015b; Woroncowa-Marcinowska et al., 2017; Hrynowiecka et al., 2018; Marciszak et al.,  
250 2019; Jach & Wojtal, 2021; Geraads et al., 2021; for more see Appendix 1) and many research  
251 projects, especially joint projects conducted by Adam Mickiewicz University, Poznań with  
252 Polish museums. These works have resulted in the discovery of a new sites, especially in  
253 Wielkopolska (Krosinko, Oborniki, Pызdry, Sławie, Turek: Pawłowska, 2009a, 2009b, 2010,  
254 2015a, 2015b, 2017b, 2022, 2023; Pawłowska et al., 2022), but also in Małopolska



255 (Siedliszowice: Pawłowska, 2012). Notable among these works is a monograph (Woroncowa-  
256 Marcinowska et al., 2013) discussing the remains of Pleistocene fauna, including fossil  
257 rhinoceroses, in their geological, historic, and chronological context, which had not been done  
258 previously on such a scale. This work also yielded an opportunity to examine the forest  
259 elephant described by Stankowski (1989), which revealed its physical condition by identifying  
260 pathologies (Pawłowska et al., forthcoming) and its depositional history from the final  
261 moments of its life (Stankowski, 1989; Pawłowska, 2009a, 2009b). Also, the find was placed  
262 among all other finds of forest elephants from Poland (Pawłowska, 2015a).

263 Recognition of Polish evidence for woolly rhinoceros' distribution was recently  
264 expanded to 157 sites by Geraads et al. (2021) who presented an update of Kowalski's  
265 synthesis, along with the revision of some site names, and demonstration of 58 new sites  
266 (Table 1). The general distributional pattern of the sites has not changed, but has densified  
267 due to the discovery of later findings by various authors (Table 1; Appendix 1).

268 A recent synthesis of fossil rhinos from Poland (Stefaniak et al., 2023) states that it  
269 "presents current state of knowledge of Quaternary rhinos from ... Poland", but also  
270 comments that "The taxonomy of rhinos was not reviewed herein, except in the case of  
271 *Stephanorhinus etruscus* and *Stephanorhinus hemitoechus*." This means that Poland still lacks  
272 a published up-to-date detailed list of woolly rhinoceros sites. Assuming that sites that are  
273 stated in that article to contain woolly rhinoceros remains actually do so, and correcting some  
274 errors in the work (there are a few in Appendix 2), it seems that Stefaniak et al. (2023) listed  
275 179 Polish sites, thus extending the list of Geraads et al. (2021), though without citing their  
276 work, with the addition of twenty two sites. Here we present a list of 215 sites from Poland  
277 where the remains of the woolly rhinoceros have been found. Thus, we are updating this list  
278 with another 36 archive sites.

279 Determining the origin of the specimens in line with the recently introduced contextual  
280 approach in palaeozoology (Pawłowska, 2022), is challenging, but will be possible after in-  
281 depth analysis. Data for Europe show that the woolly rhinoceros was widely distributed across  
282 northern Eurasia during the Pleistocene (among others: Álvarez-Lao & García, 2011;  
283 Boeskorov, 2001; Markova et al., 2013, Rey-Iglesia et al., 2021).

284

### 285 **3.2. Demographic data**

286

287 The number of identified specimens (NISP) of woolly rhinoceros from Poland is currently  
288 unknown for most sites (Table 1). Many of the assemblages have not been studied, and they  
289 particularly lack a consistent system of recording their state of preservation. One implication  
290 of this is that the number of individuals of woolly rhinoceros is also unknown. We thus still do  
291 not know how many woolly rhinoceros individuals there could have been in the Pleistocene in  
292 Poland. Individual papers have indicated the presence of single rhinoceroses (one to six: Table  
293 1 and Appendix 1), although it is difficult to determine whether the methodological approach  
294 of these assessments was the same. The minimum number of individuals of woolly rhino from  
295 Zwoleń, given by Stefaniak et al. (2023) as 20, must be wrong, since Gautier (2005) established  
296 it to be four during studying this assemblage. For Europe, the data is also sporadic and the  
297 number is usually low, not exceeding ten (four for Whitemoor Haye in United K: Schreve et  
298 al., 2013; three for Jou Puerta in Spain: Álvarez-Lao, 2014).

299 Although several works have made reference to the age of death of woolly  
300 rhinoceroses, usually using unfused epiphysis with the shaft as the criterion, the most seminal  
301 work in this regard is the monograph by Borsuk-Białynicka (1973). The age of death for Polish  
302 woolly rhinoceros individuals (and also taking into account some material from Russia and the  
303 Czech Republic) was evaluated using the skulls of the woolly rhinoceros which were assigned  
304 into age groups (Borsuk-Białynicka, 1973).

305 Gender is rarely indicated in studies of Polish woolly rhinoceros (i.a. male: Góra  
306 Puławska; Kazimierz, Konin; Józwin, Konin; Łódź; Silesia and female: Pyskowice-Rzeczyce, River  
307 Vistula (near Warsaw), NN\_4 (Borsuk-Białynicka, 1973). This may in part be due to the  
308 methodological difficulty of determining using traditional methods such as measurement.

309

### 310 **3.3. Chronological evidence**

311

312 Studying the chronology of megafauna is crucial to understanding the patterns of animal  
313 occupation of the area and the relationship of their presence with environmental change and  
314 the emergence of humans. However, in order to reconstruct these past events, it is necessary  
315 to cross-reference the results with the taphonomic events that affected the deposition of  
316 various assemblages. This is all the more important because the spatial and temporal  
317 distribution of fauna in the Pleistocene in Europe was not uniform, and some species, such as  
318 the woolly rhinoceros went extinction.

319 The remains of woolly rhinoceroses from Poland have not yet been studied with this  
320 approach. Dating of woolly rhinoceros remains has been performed by Stuart & Lister (2012)  
321 regarding the Szczecin site and Jasna Strzegowska Cave, while Wojtal (2007) has dated  
322 remains from Deszczowa Cave. The Krosinko open site is unique, and Pawłowska (2022, 2023)  
323 provided further radiocarbon dates for the woolly rhinoceros remains found there, which are  
324 embedded in the lithological and stratigraphic sequence. This is important in view of the  
325 unclear stratigraphic relations at other sites, including those from caves (or perhaps especially  
326 relating to those from caves). Moreover, the radiocarbon dates of the woolly rhinoceros from  
327 Krosinko are the oldest known from Poland (Table 2). Overall however, few radiocarbon dates  
328 are available at present — a total of  $n = 34$  from 24 sites (Table 2). When dates are excluded  
329 due to being out of range and errors (Table 2), this is reduced to 24 radiocarbon dates from  
330 17 sites. This results in the fact that there is for the time being no basis on which to infer the  
331 chronology of the woolly rhinoceros in Poland. The published dates as they are now would  
332 suggest that the remains of the woolly rhinoceros come from the range 47.3-19.5 cal. ka BP,  
333 though this does not allow use to assess the nature of the colonization of Poland by the woolly  
334 rhinoceros in the Pleistocene (Table 2). According to Marciszak et al. (2024), the youngest  
335 woolly rhinos from Poland comes from Skarszyn site, dated ca. 16.5 ka BP.

336 A detailed chronology of *C. antiquitatis* in Europe was reconstructed by Stuart & Lister  
337 (2012), with the conclusion that this species was widespread across the continent, though its  
338 range apparently contracted from ca 35 cal. ka BP. These authors' stratigraphic gap for 40–38  
339 ka BP has recently been verified (Pawłowska, 2022, 2023) by revealing the presence of woolly  
340 rhinoceros at this time (38 ka BP) in Poland, east of Krosinko, as indicated by the taphonomic  
341 data of the assemblage from this site (Pawłowska, 2023).

342

### 343 **3.4. Genetic and pathology evidences**

344

345 The maintenance of a healthy population depends on the preservation of as much genetic  
346 diversity as possible, as this affects the survival of the young, general resistance to disease,  
347 and the ability to adapt to altered conditions (Dąbrowski, 2006).

348 The usefulness of ancient DNA in inferring megafauna population sizes and diversity  
349 has been shown by many authors, such as Shapiro et al. (2004) and Campos et al. (2010), who  
350 drew conclusions regarding steppe bison and muskoxen.

351 Palaeopopulation results for woolly rhinoceroses in Poland are lacking for sufficiently  
352 large samples to allow estimation of population size and condition. Neither have pathological  
353 studies of woolly rhinoceros remains been carried out in a systematic or screening manner,  
354 which prevents inferences from being drawn concerning the condition of individuals in the  
355 Polish palaeopopulation.

356 Genetic results to date from Eastern Europe and Asia suggest that the woolly  
357 rhinoceros' population there can be considered to have been healthy, based on the lack of  
358 evidence of any decline in genetic diversity in the fossil material that has been studied from  
359 81 sites in Germany (n = 2), the North Sea (n = 6), Russia (n = 69), and China (n = 4) (Lorenzen  
360 et al., 2011).

361

### 362 **3.5. Taphonomic evidence**

363

364 Taphonomy deals with the transition of animal remains from the biosphere to the lithosphere  
365 (Efremov, 1940) and involves all processes that act on organic remains, from the death of the  
366 organism through fossilization (Behrensmeyer & Hill, 1980). These processes affect the degree  
367 and quality of preservation of animal remains. The use of taphonomic analysis in studies of  
368 Pleistocene faunal material provides an opportunity to assess the factors (including cultural  
369 factors) responsible for the modifications, reworking, and accumulation of material at a given  
370 site (Pawłowska, 2010, 2023).

371 Although the general suitability of taphonomic analysis for palaeozoological research  
372 of Polish assemblages has already been shown (Pawłowska, 2010), it is relatively rarely  
373 employed in these studies. There are several reasons for this: (1) the poor preservation of the  
374 material may make it difficult to carry out taphonomic studies or to recognize marks; (2)  
375 taphonomy requires knowledge of the scope of the marks and an ability to recognise them;  
376 and (3) taphonomy requires high-resolution methods. Moreover, the limitations seen to date  
377 on the use of taphonomic analysis in the study of Polish Pleistocene faunal material from caves  
378 shows that it is necessary to undertake such studies on materials from a fluvial context —  
379 which are the most numerous materials represented in Polish Pleistocene faunal assemblages  
380 — and from eolian contexts.

381 To date, in studies of Polish collections, taphonomic studies of fossil rhinoceroses have  
382 been carried out in the fluvial context for both the forest rhino (*Stephanorhinus*

383 *kirchbergensis*; known from 14 sites in Poland) (Geraads et al., 2021; Pawłowska, 2017  
384 unpublished results;) and the woolly rhino (*Coelodonta antiquitatis*) (Pawłowska, 2023). In the  
385 case of the forest rhino (*Stephanorhinus kirchbergensis*) from Gorzów Wielkopolski, the  
386 taphonomic study did not reveal marks resulting from hominid activity. Instead, these studies  
387 made it possible to discover and study by Kamilla Pawłowska pathological changes, which  
388 were located mainly in the joints (Pawłowska, 2017 unpublished results). In the case of the  
389 woolly rhinoceros, taphonomic analysis revealed a secondary context for its remains, in the  
390 form of the effects of a braided river as depositional factor, and showed, for the first time in  
391 a Polish study, the effect of temperature on rhino bones, in the form of burn marks  
392 (Pawłowska, 2023).

393 For other assemblages from Poland, evidence of processing in the form of potential  
394 cultural marks needs to be verified using modern techniques. Cut marks on a woolly  
395 rhinoceros mandible were described by Bratlund (2005) as evidence of defleshing of the head  
396 in order to extract the muscle. Other marks identified as cut marks on animal bones from this  
397 site were however deemed not to be evidence of human activity, and are instead regarded as  
398 resulting from natural causes, such as trampling. They are thus not cultural marks (Gautier,  
399 2005).

400 Knowledge of cultural marks from the Pleistocene is very relevant to identifying the  
401 activity associated with the type of marks, and thus in understanding whether access to  
402 carcasses was early or late, the stages of the processing of carcasses, and the manufacturing  
403 process. These topics have been treated by various authors, who have formulated a  
404 palaeogeography of butchering for Pleistocene (Pawłowska, 2017a) and subsistence strategy  
405 (Bunn, 1986; Alhaique et al., 2004; Rivals et al., 2006; Domínguez-Rodrigo et al., 2015; Van  
406 Kolfschoten et al., 2015; Espigares et al., 2019).

407 Social environment can also be studied by other marks induced by hominids  
408 (Pawłowska, 2020), such as worked pieces. However, there is a little data in this respect from  
409 Poland (Pawłowska, 2022).

410 Our summary of studies of Polish woolly rhinoceroses has also shown that none of the  
411 studies conducted to date have included taphonomic analysis on a countrywide scale in order  
412 to identify factors of faunal material deposition. Individual works have drawn attention to  
413 marks of biting by predators, the influence of water, and other effects (Appendix 1).

414

415 **3.6. Art**

416

417 There is no evidence of Palaeolithic art or worked objects involving the woolly rhinoceros from  
418 Poland.

419

420 **3.7. Summary of state of art and perspectives**

421

422 The woolly rhinoceros (*Coelodonta antiquitatis* Blumenbach 1799) and the woolly mammoth  
423 are the main representatives of the Pleistocene megafauna. To date, however, the majority  
424 of scholarly attention has been paid to the woolly mammoth, the remains of which have  
425 received morphometric, radiocarbon dating, DNA, and isotope analysis. There is a surprising  
426 dearth of research on the woolly rhinoceros given the relative abundance of its remains in  
427 Poland (Kowalski, 1959; Geraads et al., 2021). We thus have no knowledge of general  
428 demographic data, phylogenetic relationships, or other details of the presence of woolly  
429 rhinoceroses in Poland. For example, we do not know if the woolly rhinoceros' presence was  
430 permanent, temporary, or periodic. We also lack knowledge in relation to other aspects —  
431 such as, for example, the condition of the animals, which would assist in reconstructing the  
432 life and death of individuals. The archive data remains scattered, but research and metadata  
433 collection should make it possible to create a comprehensive database and to draw useful  
434 conclusions.

435 To address this gap, a multifaceted study, the WOOLRHINOPOLI Project, was proposed  
436 in 2020, and since 2022 has been implemented step by step in line with the issues to be  
437 examined and the work schedules. The WOOLRHINOPOLI Project brings together eleven  
438 researchers from Europe with the aim of unravelling the chronological, geographical, and  
439 taphonomic complexities of the occurrence of the woolly rhinoceros in the Pleistocene  
440 contexts of Poland (WOOLRHINOPOLI) and Europe. Since reliable conclusions can only be  
441 drawn on the basis of the metadata, and not from studies that deal with individual samples,  
442 reaching these milestones on the European scale will involve examining the remains of woolly  
443 rhinoceroses from Poland, the North Sea, and selected European countries (Germany, the  
444 Czech Republic, the Netherlands, the United Kingdom, France, Spain, Italy, Romania, Beringia,  
445 and Moldova).

446

#### 447 4. Conclusions

448

449 The woolly rhinoceros, an extinct representative of the megafauna, has been the subject of  
450 Polish research since the nineteenth century. Our focus has been on its remains found in  
451 various archive locations in Poland, summarizing our knowledge of this taxon as a starting  
452 point for further multifaceted research that will be implemented. The conclusions of previous  
453 research are as follows:

- 454 • The lack of verification of the taxonomic and anatomical designations of woolly rhinoceros  
455 remains since the 1960s, as well as the lack of large-scale field research, means that the  
456 quantity of woolly rhinoceros remains, the number of individuals, and consequently our  
457 state of knowledge of sites with woolly rhinoceros remains in Poland are unknown.
- 458 • As a consequence of the first point, details of the distribution of the woolly rhinoceros in  
459 Poland, as well as in Central and Western Europe, are currently unknown.
- 460 • The limited number of currently available published radiocarbon dates for the remains of  
461 the woolly rhinoceros does not allow for the reconstruction of its chronology in Poland  
462 during the Pleistocene and Holocene. Also, in light of the development of research  
463 methods, the radiocarbon dates alone are insufficient to provide a scenario of events of  
464 the occupation of Poland by the woolly rhinoceros.
- 465 • The analysis of taphonomical characteristics from the biostratigraphical and diagenetic  
466 stages is a necessary approach to studies of Pleistocene mammals, and such studies should  
467 be deepened and more widely applied to woolly rhinoceros remains.
- 468 • Several issues regarding hominid–rhinoceros relationships have been addressed for  
469 individual sites in Polish studies (Pawłowska, 2022, 2023); this also sets the direction for  
470 further in-depth and synthetic analyses.
- 471 • There is no known figural depictions or paintings showing depictions of the woolly  
472 rhinoceros from Poland.
- 473 • The state of research on woolly rhinoceros remains is, unfortunately, unsatisfactory, but  
474 this allows us to set the direction for the further research currently being carried out under  
475 the auspices of the WOOLRHINOPOLI Project.
- 476 • The main conclusion is thus that the woolly rhinoceros, as an extinct species of rhino and  
477 the main representative of the *Mammuthus–Coelodonta* Faunal Complex (Kahlke, 2014),

478 deserves more attention and systematic research in Polish research that aims to reveal its  
479 Quaternary history.

480

## 481 **Acknowledgments**

482 This research was funded in whole or in part by the National Science Center, Poland  
483 (2021/43/B/ST10/00362; WOOLRHINOPOLI). For the purpose of Open Access, the author has  
484 applied a CC-BY public copyright license to any Author Accepted Manuscript (AAM) version  
485 arising from this submission.

486 We thank two anonymous reviewers for their comments and suggestions, which  
487 contributed to improving the content of this paper.

## 488 **Appendices**

489 Appendix 1 and 2 are available on <http://www.geologos.com.pl/>

490 Appendix 1. List of publications on the study of fossil rhinoceros remains from Poland  
491 discusses here, as of February 2024.

492 Appendix 2. A list of changes made to the scope of the recent synthesis (Stefaniak et al., 2023)  
493 of fossil rhinoceros remains, including woolly rhinoceros, which were necessary for the  
494 compilation presented here.

495

## 496 **References**

- 497 Alhaique, F., Bisconti, M., Castiglioni, E., Cilli, C., Fasani, L., Giacobini, G., Grifoni, R.,  
498 Guerreschi, A., Iacopini, A., Malerba, G., Peretto, C., Recchi, A., Ris, A.R., Ronchitelli, A.,  
499 Rottoli, M., Hohenstein, U.T., Tozzi, C., Visentini, P. & Wilkens, B., 2004. Animal resources  
500 and subsistence strategies. *Collegium Antropologicum* 28, 23–40.
- 501 Álvarez-Lao, D.J., 2014. New discoveries of woolly mammoth and woolly rhinoceros from  
502 Northern Iberia. In Abstract book of the VIth International Conference on Mammoths  
503 and their relatives. SASG, Special, 102, 19.
- 504 Álvarez-Lao, D.J. & García, N., 2011. Southern dispersal and Palaeoecological implications of  
505 woolly rhinoceros (*Coelodonta antiquitatis*): review of the Iberian occurrences.  
506 *Quaternary Science Reviews* 30, 2002–2017.



507 Ballatore, M., 2016. Palaeoecological investigations on Plio-Pleistocene European  
508 Rhinoceroses (Genus *Stephanorhinus*): powder X-ray diffraction, carbone isotope  
509 geochemistry, tooth wear analyses and biometry. *Plinius* 42, 16–19.

510 Ballatore, M. & Breda, M., 2019. Revision of the rhinoceros remains (Rhinocerotidae,  
511 Mammalia) from the late Pliocene of Etouaires (Auvergne, France) and the morphological  
512 distinction between the postcranial bones of *Stephanorhinus elatus* and *S. etruscus*.  
513 *Comptes Rendus Palevol* 18, 191–208.

514 Behrensmeyer, A.K. & Hill, A.P. (Eds), 1980. *Fossils in the making: vertebrate taphonomy and*  
515 *paleoecology* (No. 69). University of Chicago Press. 345 pp.

516 Belyaev, R.I., Boeskorov, G.G., Cheprasov, M.Y. & Prilepskaya, N.E., 2023. A new discovery in  
517 the permafrost of Yakutia sheds light on the nasal horn morphology of the woolly  
518 rhinoceros. *Journal of Morphology* 284, e21626.

519 Boeskorov, G., 2001. Woolly rhino (*Coelodonta antiquitatis*) distribution in Northeast Asia.  
520 *Deinsea* 8, 15–20.

521 Boeskorov, G.G., Lazarev, P.A., Sher, A.V., Davydov, S.P., Bakulina, N.T., Shchelchkova, M.V.,  
522 Binladen, J., Willerslev, E., Buigues, B. & Tikhonov, A.N., 2011. Woolly rhino discovery in  
523 the lower Kolyma River. *Quaternary Science Reviews* 30, 2262–2272.

524 Borsuk-Białynicka, M., 1973. Studies on the Pleistocene rhinoceros *Coelodonta antiquitatis*  
525 (Blumenbach). *Palaeontologia Polonica* 29, 1–152.

526 Bratlund, B., 2005. Comment on a cut-marked woolly rhino mandible from Zwoleń. [In:] R.  
527 Schild (Ed.): *The killing fields of Zwoleń. A Middle Paleolithic kill-butcher-site in Central*  
528 *Poland*. Institute of Archaeology and Ethnology, Polish Academy of Sciences, Warsaw,  
529 217–221.

530 Bunn, H.T., 1986. Patterns of skeletal representation and hominid subsistence activities at  
531 Olduvai Gorge, Tanzania, and Koobi Fora, Kenya. *Journal of Human Evolution* 15, 673–690.

532 Campos, P.F., Willerslev, E., Sher, A., Orlando, L., Axelsson, E., Tikhonov, A., Aaris-Sørensen,  
533 K., Greenwood, A.D., Kahlke, R.-D., Kosintsev, P., Krakhmalnaya, T., Kuznetsova, T., Lemey,  
534 P., MacPhee, R., Norris, C.A., Shepherd, K., Suchard, M.A., Zazula, G.D., Shapiro, B. &  
535 Gilbert, M.T.P., 2010. Ancient DNA analyses exclude humans as the driving force behind  
536 late Pleistocene musk ox (*Ovibos moschatus*) population dynamics. *Proceedings of the*  
537 *National Academy of Sciences* 107, 5675–5680.

538 Dąbrowski, I., 2006. White rhinoceros, *Ceratotherium simum* (Burchell, 1817) and black  
539 rhinoceros, *Diceros bicornis* (Linne, 1758) breeding problems in vivarium conditions.  
540 Warszawa. 124 pp.

541 Deng, T., Wang, X., Fortelius, M., Li, Q., Wang, Y., Tseng, Z.J., Takeuchi, G.T., Saylor, J.E., Saila,  
542 L.K. & Xie, G., 2011. Out of Tibet: Pliocene woolly rhino suggests high-plateau origin of Ice  
543 Age megaherbivores. *Science* 333 (6047), 1285–1288.

544 Domínguez-Rodrigo, M., Barba, R., Soto, E., Sesé, C., Santonja, M., Pérez-González, A.,  
545 Yravedra, J. & Galán, A.B., 2015. Another window to the subsistence of Middle Pleistocene  
546 hominins in Europe: A taphonomic study of Cuesta de la Bajada (Teruel, Spain).  
547 *Quaternary Science Reviews* 126, 67–95.

548 Efremov, J.A., 1940. Taphonomy: a new branch of geology. *Pan-American Geologist* 74, 81–  
549 93.

550 Espigares, M. P., Palmqvist, P., Guerra-Merchán, A., Ros-Montoya, S., García-Aguilar, J.M.,  
551 Rodríguez-Gómez, G., Serrano, F.J. & Martínez-Navarro, B., 2019. The earliest cut marks  
552 of Europe: a discussion on hominin subsistence patterns in the Orce sites (Baza Basin, SE  
553 Spain). *Scientific reports* 9, 1–13.

554 Fortelius, M., 1983. The morphology and palaeobiological significance of the horns of  
555 *Coelodonta antiquitatis* (Mammalia: Rhinocerotidae). *Journal of Vertebrate Paleontology*  
556 3, 125–135.

557 García-Fernández, D., Cerdeño, E., Sanz, M. & Daura, J., 2023. The Latest Occurrence of  
558 *Stephanorhinus hundsheimensis* (Rhinocerotidae) in Europe: The Skeletons from the Cova  
559 del Rinoceront Site (Castelldefels, Barcelona). *Quaternary* 6, 60.

560 Gautier, A., 2005. The Zwoleń mammals. [In:] R. Schild (Ed): *The killing fields of Zwoleń: a*  
561 *middle Paleolithic kill-butcher-site in Central Poland*. Polish Academy of Sciences.  
562 Institute of Archaeology and Ethnology, 71–109.

563 Geraads, D., Cerdeño, E., García Fernandez, D., Pandolfi, L., Billia, E., Athanassiou, A., Albayrak,  
564 E., Codrea, V., Obada, T., Deng, T., Tong, H., Lu, X., Pícha, S., Marciszak, A., Jovanovic, G.,  
565 Becker, D., Zervanova, J., Chaïd Saoudi, Y., Bacon, A.-M., Sévêque, N., Patnaik, R., Brezina,  
566 J., Spassov, N. & Uzunidis, A., 2021. A Database of Old World Neogene and Quaternary  
567 Rhino-bearing Localities. Available at [www.rhinosourcecenter.com/about/fossil-rhino-](http://www.rhinosourcecenter.com/about/fossil-rhino-database.php)  
568 [database.php](http://www.rhinosourcecenter.com/about/fossil-rhino-database.php).

569 Hrynowiecka, A., Żarski, M., Jakubowski, G., Nadachowski, A., Pawłowska, K., Pawłowski, D.,  
570 Szymanek, M. & Nast, D., 2018. Eemian and Vistulian (Weichselian) paleoenvironmental  
571 changes: A multi-proxy study of sediments and mammal remains from the Ławy paleolake  
572 (Eastern Poland). *Quaternary International* 467, 131–146.

573 Hrynowiecka, A., Żarski, M., Chmielowska, D., Pawłowska, K., Okupny, D., Michczyński, A. &  
574 Kukulak, J., 2022. Reconstruction of 26 kyrs palaeoenvironmental history of the Czarny  
575 Dunajec Fan—A multiproxy study of the Długopole gravel pit deposits (Western  
576 Carpathians, S Poland). *Catena* 211, 105940.

577 Jach, R. & Wojtal, P., 2021. Taphonomy of Pleistocene large mammal remains in the deposits  
578 of river Raba, southern Poland. *Annales Societatis Geologorum Poloniae* 91, 167–187.

579 Kaczmarek, J. 2004. Kości zostały rzucone. W [ ]: Perty z lamusa: Archeologia w dokumentach  
580 Archiwum Naukowego Muzeum Archeologicznego w Poznaniu, 1–3.

581 Kahlke, R.D., 2014. The origin of Eurasian mammoth faunas (*Mammuthus–Coelodonta* faunal  
582 complex). *Quaternary Science Reviews* 96, 32–49.

583 Kahlke, R.D. & Lacomat, F., 2008. The earliest immigration of woolly rhinoceros (*Coelodonta*  
584 *tologojensis*, Rhinocerotidae, Mammalia) into Europe and its adaptive evolution in  
585 Palaeartic cold stage mammal faunas. *Quaternary Science Reviews* 27, 1951–1961.

586 Karaszewski, W., 1980. Szczątki kostne nosorożca włochatego (*Coelodonta antiquitatis*  
587 Blumenbach) w Przasnyszu [Bone remains of the woolly rhinoceros (*Coelodonta*  
588 *antiquitatis* Blumenbach) in Przasnysz]. *Kwartalnik Geologiczny* 24, 857–859.

589 Kirillova, I.V., Chernova, O.F., Van Der Made, J., Kukarskih, V.V., Shapiro, B., Van Der Plicht, J.,  
590 Shidlovskiy, F.K., Heintzman, P.D., Van Kolfschoten, T. & Zaninai, O.G., 2017. Discovery of  
591 the skull of *Stephanorhinus kirchbergensis* (Jäger, 1839) above the Arctic Circle.  
592 *Quaternary Research* 88, 537–550.

593 Kowalski, K., 1959. Katalog ssaków plejstocenu Polski [A catalogue of the Pleistocene  
594 mammals of Poland]. Państwowe Wydawnictwo Naukowe, Warszawa and Wrocław, 267  
595 pp.

596 Kowalski, K., 2000. Der pleistozäne Ölsumpf bei Starunia, Ukraine. *Europäische*  
597 *Fossilagerstätten*, 232–236.

598 Król, P., 1998. Ssaki kopalne czwartorzędu Polski w zbiorach przyrodniczych Muzeum  
599 Narodowego w Kielcach [Fossil mammals of the Quaternary of Poland in the natural

600 history collection of the National Museum in Kielce]. *Rocznik Muzeum Narodowego w*  
601 *Kielcach* 19, 309–316.

602 Kubiak, H., 1969. Über die Bedeutung der Kadaver des Wollhaarnashorns von Starunia.  
603 *Berichte der Deutschen Gesellschaft für Geologische Wissenschaften Series A Geologie und*  
604 *Palaontologie* 14, 345–347.

605 Kubiak, H., 1989. Nieparzystokopytne-Perissodactyla [Equidae - Perissodactyla]. *Folia*  
606 *Quaternaria* 59-60, 197–201.

607 Kubiak, H., 1994. Saturnia w 85 rocznicę pierwszych odkryć paleontologicznych [Saturnia on  
608 the 85th anniversary of the first paleontological discoveries]. *Wszecławiat* 95, 295–299.

609 Kubiak, H., 2009. The Starunia collections in the Institute of Systematics and Evolution of  
610 Animals, Polish Academy of Sciences, Krakow. *Geoturystyka* 18, 71–80.

611 Kubiak, H. & Drygant, D.M., 2005. The Starunia collections in Lviv and Krakow Natural History  
612 Museums and history of paleontological studies. [In:] M.J. Kotarba (Ed.): *Geological*  
613 *Studies 2004-2005 at Starunia – The area of discoveries of woolly rhinoceroses*. Warsaw-  
614 Krakow, Polish Geological Institute and Society of Research on Environmental Changes  
615 “Geosphere”, 37–44.

616 Kubiak, H. & Dziurdzik, B., 1973. Histological characters of hairs in extant and fossil  
617 rhinoceroses. *Acta Biologica Cracoviensia* 16, 55–61.

618 Kuc, T., Różański, K., Goslar, T., Kubiak, H. & Kotarba, M.J., 2005. Radiocarbon dating of  
619 remnants of Woolly Rhinoceroses and Mammoth from Starunia, fore-Carpathian region,  
620 Ukraine. [In:] M.J. Kotarba (Ed): *Geological Studies 2004-2005 at Starunia – The area of*  
621 *discoveries of woolly rhinoceroses*. Warsaw-Krakow, Polish Geological Institute and  
622 Society of Research on Environmental Changes “Geosphere”, 195–203.

623 Kuc, T. Różański, K., Kotarba, M.J., Goslar, T. & Kubiak, H., 2012. Radiocarbon dating of  
624 Pleistocene Fauna and Flora from Starunia, SW Ukraine. *Radiocarbon* 54, 123–126.

625 Lord, E., Dussex, N., Kierczak, M., Díez-del-Molino, D. & Ryder, O.A., 2020. Pre-extinction  
626 Demographic Stability and Genomic Signatures of Adaptation in the Woolly Rhinoceros.  
627 *Current Biology* 30, 3871–3879.

628 Lorenzen, E.D., Nogués-Bravo, D., Orlando, L., Weinstock, J., Binladen, J., Marske, K.A., Ugan,  
629 A., Borregaard, M., Gilbert, T.P., Nielsen, R., Ho, S.Y.W., Goebel, T., Graf, K.E., Byers, D.,  
630 Stenderup, J.T., Rasmussen, M., Campos, P.F., Leonard, J.A., Koepfli, K-P., Froese, D.,  
631 Zazula, G., Stafford, T.W., Aaris-Sørensen, K., Batra, P., Haywood, A.M., Singarayer, J.S.,

632 Valdes, P.J., Boeskorov, G., Burns, J.A., Davydov, S.P., Haile, J., Jenkins, D.L., Kosintsev, P.,  
633 Kuznetsova, T., Lai, X., Martin, L.D., McDonald, H.G., Mol, D., Meldgaard, M., Munch, K.,  
634 Stephan, E., Sablin, M., Sommer, R.S., Sipko, T., Scott, E., Suchard, M.A., Tikhonov, A.,  
635 Willerslev, R., Wayne, R.K., Cooper, A., Hofreiter, M., Sher, A., Shapiro, B., Rahbek, C. &  
636 Willerslev, E., 2011. Species-specific responses of Late Quaternary megafauna to climate  
637 and humans. *Nature* 479, 359–364.

638 Lubicz-Niezabitowski, E., 1913. Czaszka nosorożca włochatego (*Rhinoceros antiquitatis* Blum.)  
639 ze Złotego Potoka [Skull of a woolly rhinoceros (*Rhinoceros antiquitatis* Blum.) from Złoty  
640 Potok]. *Sprawozdanie Komisji Fizyograficznej PAU w Krakowie* 47, 3–11 (in Polish with  
641 German summary).

642 Lubicz-Niezabitowski, E., 1926. Szczątki nosorożca włochatego (*Rhinoceros antiquitatis* Blum.)  
643 znalezione na ziemi wielkopolskiej [Remains of a woolly rhinoceros (*Rhinoceros*  
644 *antiquitatis* Blum.) found in Greater Poland]. *Muzeum Wielkopolskie w Poznaniu* 2, 1–52  
645 (in Polish with French summary).

646 Lubicz-Niezabitowski, E., 1929. Interglacjał w Szelażu pod Poznaniem (Interglazial in Szelaż  
647 [Schilling] bei Posen). In: Część II. Fauna pokładów drugiego okresu międzylodowcowego  
648 w Szelażu, *Sprawozdanie Komisji Fizjograficznej PAU* 63, 51–70 (in Polish with German  
649 summary).

650 Lubicz-Niezabitowski, E., 1938. O kilku ciekawszych szczątkach kopalnych zwierząt ssących  
651 Polski [About some of the more interesting fossil remains of Polish mammals].  
652 *Sprawozdania Poznańskiego Towarzystwa Przyjaciół Nauk* 11, 77–78 (in Polish).

653 Marciszak, A., Kotowski, A., Przybylski, B., Badura, J., Wiśniewski, A. & Stefaniak, K., 2019.  
654 Large mammals from historical collections of open-air sites of Silesia (southern Poland)  
655 with special reference to carnivores and rhinoceros. *Historical Biology* 31, 696–730.

656 Markova, A. K., Puzachenko, A.Y., Van Kolfschoten, T., Van der Plicht, J. & Ponomarev, D.V.,  
657 2013. New data on changes in the European distribution of the mammoth and the woolly  
658 rhinoceros during the second half of the Late Pleistocene and the early Holocene.  
659 *Quaternary International* 292, 4–14.

660 Orlando, L., Leonard, J. A., Thenot, A., Laudet, V., Guerin, C. & Hänni, C., 2003. Ancient DNA  
661 analysis reveals woolly rhino evolutionary relationships. *Molecular phylogenetics and*  
662 *evolution* 28, 485–499.

- 663 Pandolfi, L., 2023. Reassessing the phylogeny of Quaternary Eurasian Rhinocerotidae. *Journal*  
664 *of Quaternary Science* 38, 291–294.
- 665 Pandolfi, L., Boscato, P., Crezzini, J., Gatta, M., Moroni, A., Rolfo, M. & Tagliacozzo, A., 2017.  
666 Late Pleistocene last occurrences of the narrow-nosed rhinoceros *Stephanorhinus*  
667 *hemitoechus* (Mammalia, Perissodactyla) in Italy. *Research in Paleontology and*  
668 *Stratigraphy* 123, 177–192.
- 669 Pandolfi, L., Cerdeño, E., Codrea, V. & Kotsakis, T., 2017. Biogeography and chronology of the  
670 Eurasian extinct rhinoceros *Stephanorhinus etruscus* (Mammalia, Rhinocerotidae).  
671 *Comptes Rendus Palevol* 16, 762–773.
- 672 Pawłowska, K., 2009a. Plejstocenińska bioróżnorodność faunistyczna w Krosinku [Pleistocene  
673 faunal biodiversity in Krosinko]. [In:] B. Walna, I. Kaczmarek, M. Lorenc & R. Dondajewska  
674 (Eds): *Wielkopolski Park Narodowy w badaniach przyrodniczych*. Poznań–Jeziory, 207–  
675 213.
- 676 Pawłowska, K., 2009b. Tafonomia plejstocenińskich kości ssaków [Taphonomy of Pleistocene  
677 mammalian bones]. [In:] M. Żarski & S. Lisicki (Eds): *Strefa marginalna lądolodu*  
678 *złodowacenia warty i pojezierza plejstocenińskie na południowym Podlasiu*. Warszawa, PIG-  
679 PIB: 78–80.
- 680 Pawłowska, K., 2010. The usefulness of a taphonomic approach for studies of Pleistocene  
681 mammals. *Geologos* 16, 183–189.
- 682 Pawłowska, K., 2012. Study of the Pleistocene fauna from the Tarnów area. Archive Work, 1–  
683 2.
- 684 Pawłowska, K., 2015a. Elephantids from Pleistocene Poland: State of knowledge. *Quaternary*  
685 *International* 379, 89–105.
- 686 Pawłowska, K., 2015b. Studies on Pleistocene and Holocene mammals from Poland: the legacy  
687 of Edward Feliks Lubicz-Niezabitowski (1875–1946). *Quaternary International* 379, 118–  
688 127.
- 689 Pawłowska, K., 2017a. Large mammals affected by hominins: Paleogeography of butchering  
690 for the European Early and Middle Pleistocene. *Quaternary International* 438, 104–115.
- 691 Pawłowska, K., 2017b. Taxonomic report of the analysis of specimens from the collection of  
692 the Archaeological Museum in Poznań. Archive work 1–2.
- 693 Pawłowska, K., 2020. Time of change: cattle in the social practices of Late Neolithic  
694 Çatalhöyük. *Archaeological and Anthropological Sciences* 12, 1–18.

695 Pawłowska, K., 2022. MIS 3–1 fauna from Krosinko: Implications for the past biogeography,  
696 chronology and palaeoenvironments of Poland. *Quaternary International* 632, 79–93.

697 Pawłowska, K., 2023. In front of the retreating ice-sheet: Fauna complex of central-western  
698 Poland in MIS 3–2 (Krosinko site). *Quaternary International* 674–675, 138–151.

699 Pawłowska, K., Greenfield, H. & Czubla, P., 2014. “Steppe” mammoth (*Mammuthus*  
700 *trogontherii*) remains in their geological and cultural context from Bełchatów (Poland): A  
701 consideration of human exploitation in the Middle Pleistocene. *Quaternary International*  
702 326–327, 448–468.

703 Pawłowska, K., Zieliński, T., Woronko, B., Sobkowiak-Tabaka, I. & Stachowicz-Rybka, R., 2022.  
704 Integrated environmental records in Late Pleistocene Poland: The paleofluvial regime and  
705 paleoclimate inferred from Krosinko site. *Quaternary International* 616, 12–29.

706 Prothero, D.R., 1993. Fifty million years of rhinoceros evolution. [In:] O.A. Ryder (Ed.):  
707 *Rhinoceros Biology And Conservation*. Zoological Society of San Diego, 82–91.

708 Pushkina, D., Saarinen J., Ziegler, R. & Bocherens, H., 2020. Stable isotopic and mesowear  
709 reconstructions of paleodiet and habitat of the Middle and Late Pleistocene mammals in  
710 south-western Germany. *Quaternary Science Reviews* 227, 106026.

711 Puzachenko, A.Y., Markova, A.K. & Pawłowska, K., 2022. Evolution of Central European  
712 regional mammal assemblages between the late Middle Pleistocene and the Holocene  
713 (MIS7–MIS1). *Quaternary International* 633, 80–102.

714 Rey-Iglesia, A., Lister, A.M., Stuart, A.J., Bocherens, H., Szpak, P., Willerslev, E. & Lorenzen,  
715 E.D., 2021. Late Pleistocene paleoecology and phylogeography of woolly rhinoceroses.  
716 *Quaternary Science Reviews* 263, 106993.

717 Rivals, F., Testu, A., Moigne, A.M. & de Lumley, H., 2006. The Middle Pleistocene argali (*Ovis*  
718 *ammon antiqua*) assemblages at the Caune de l’Arago (Tautavel, Pyrénées-Orientales,  
719 France): were prehistoric hunters or carnivores responsible for their accumulation?.  
720 *International Journal of Osteoarchaeology* 16, 249–268.

721 Schreve, D., Howard, A., Currant, A., Brooks, S., Buteux, S., Coope, R., Crocker, B., Field, M.,  
722 Greenwood, M., Greig, J. & Toms, P., 2013. A Middle Devensian woolly rhinoceros  
723 (*Coelodonta antiquitatis*) from Whitemoor Haye Quarry, Staffordshire (UK):  
724 palaeoenvironmental context and significance. *Journal of Quaternary Science* 28, 118–  
725 130.

726 Shapiro, B., Drummond, A.J., Rambaut, A., Wilson, M.C., Matheus, P.E., Sher, A.V., Pybus, O.G.,  
727 Gilbert, M.T.P., Barnes, I., Binladen, J., Willerslev, E., Hansen, A.J., Baryshnikov, G.F.,  
728 Burns, J.A., Davydov, S., Driver, J.C., Froese, D.G., Harington, C.R., Keddie, G., Kosintsev,  
729 P., Kunz, M.L., Martin, L.D., Stephenson, R.O., Storer, J., Tedford, R., Zimov, S. & Cooper,  
730 A., 2004. Rise and fall of the Beringian steppe bison. *Science* 306, 1561–1565.

731 Shpansky, A.V., 2017. Questions paleozoogeography of *Stephanorhinus kirchbergensis* (Jäger,  
732 1839) (Rhinocerotidae, Mammalia). *Geosphere Research* 3, 74–89.

733 Stankowski, W., 1989. Pozycja stratygraficzna szkieletu słonia leśnego w świetle analizy  
734 geologicznej, stanowisko Joźwin 1984 (doniesienie wstępne) [Stratigraphic position of the  
735 forest elephant skeleton in the light of geological analysis, site Joźwin 1984 (preliminary  
736 report)]. *Zeszyty Muzealne* 2, 95–102.

737 Staszic, S., 1815. *O ziemiordztwie Karpatów i innych gór i równin Polski [On the land of the*  
738 *Carpathians and other mountains and plains of Poland]*. Drukarnia Rządowa, Warszawa,  
739 390 pp.

740 Stefaniak, K., Stachowicz-Rybka, R., Borówka, R. K., Hrynowiecka, A., Sobczyk, A., Moskal-del  
741 Hoyo, M., Kotowski, A., Nowakowski, D., Krajcarz, M.T., Billia, E.M.E., Persico, D.,  
742 Burkanova, E.M., Leshchinskiy, S.V., van Asperen, E., Ratajczak, U., Shpansky, A.V.,  
743 Lempart, M., Wach, B., Niska, M., van der Made, J. & Kovalchuk, O., 2021. Browsers,  
744 grazers or mix-feeders? Study of the diet of extinct Pleistocene Eurasian forest rhinoceros  
745 *Stephanorhinus kirchbergensis* (Jäger, 1839) and woolly rhinoceros *Coelodonta*  
746 *antiquitatis* (Blumenbach, 1799). *Quaternary International* 605, 192–212.

747 Stefaniak, K., Kovalchuk, O., Ratajczak-Skrzatek, U., Kropczyk, A., Mackiewicz, P., Kłys, G.,  
748 Krajcarz, M., Krajcarz, M.T., Nadachowski, A., Lipecki, G., Karbowski, K., Ridush, B., Sabol,  
749 M. & Płonka, T., 2023. Chronology and distribution of Central and Eastern European  
750 Pleistocene rhinoceroses (Perissodactyla, Rhinocerotidae) - A review. *Quaternary*  
751 *International* 674-375, 87–108.

752 Stuart, A.J. & Lister, A.M., 2012. Extinction chronology of the woolly rhinoceros *Coelodonta*  
753 *antiquitatis* in the context of late Quaternary megafaunal extinctions in northern Eurasia.  
754 *Quaternary Science Reviews* 51, 1–17.

755 Tiunov, A.V. & Kirillova, I.V., 2010. Stable isotope ( $^{13}\text{C}/^{12}\text{C}$  and  $^{15}\text{N}/^{14}\text{N}$ ) composition of the  
756 woolly rhinoceros *Coelodonta antiquitatis* horn suggests seasonal changes in the diet.  
757 *Rapid Communications in Mass Spectrometry* 24, 3146–3150.



758 Uzunidis, A., Antoine, P.O. & Brugal, J.P., 2022. A Middle Pleistocene *Coelodonta antiquitatis*  
759 *praecursor* (Mammalia, Perissodactyla) from Les Rameaux, SW France, and a revised  
760 phylogeny of *Coelodonta*. *Quaternary Science Reviews* 288, 107594.

761 van Asperen, E.N. & Kahlke, R.D., 2015. Dietary variation and overlap in Central and Northwest  
762 European *Stephanorhinus kirchbergensis* and *S. hemitoechus* (Rhinocerotidae,  
763 Mammalia) influenced by habitat diversity: "You'll have to take pot luck!" (proverb).  
764 *Quaternary Science Reviews* 107, 47–61.

765 van Kolfschoten, T., Buhrs, E. & Verheijen, I., 2015. The larger mammal fauna from the Lower  
766 Paleolithic Schöningen Spear site and its contribution to hominin subsistence. *Journal of*  
767 *Human Evolution* 89, 138–153.

768 Vereshchagin, N. K. & Baryshnikov, G.F., 1992. The ecological structure of the "Mammoth  
769 Fauna" in Eurasia. *Annales Zoologici Fennici* 28, 253–259.

770 Wiszniowska, T., Stefaniak, K. & Socha, P., 2005. Szczątki kręgowców ze stanowiska środkowo  
771 paleolitycznego przy ul. Hallera we Wrocławiu [Vertebrate remains from the Middle  
772 Paleolithic site at ul. Hallera in Wrocław]. *Śląskie Sprawozdania Archeologiczne* 47, 17–  
773 23.

774 Wojtal, P., 2007. *Zooarcheological studies of the Late Pleistocene sites in Poland*. Institute of  
775 Systematics and Evolution of Animals, Polish Academy of Sciences, Kraków, 189 pp.

776 Wolfhagen, J., Veropoulidou, R., Ayala, G., Filipović, D., Kabukcu, K., Lancelotti, C., Madella,  
777 M., Pawłowska, K., Santiago-Marrero, C.G. & Wainwright, J., 2020. The Seasonality of  
778 Wetland and Riparian Taskscapes at Çatalhöyük. *Near Eastern Archaeology* 83, 98–109.

779 Woroncowa-Marcinowska, T., Żarski, M., Pawłowska, K. & Urban, J., 2013. *Rewizja i naukowe*  
780 *opracowanie kości ssaków zgromadzonych w kolekcjach Muzeum Geologicznego PIG-PIB*  
781 *[Revision and scientific analysis of mammal bones collected in the collections of the PGI-*  
782 *PIB Geological Museum]*. Narodowe Archiwum Geologiczne, 130 pp.

783 Woroncowa-Marcinowska, T., Pawłowska, K., Żarski, M. & Urban, J., 2017. Zespoły  
784 plejstocenijskiej fauny (zbiory Muzeum Geologicznego PIG-PIB) w ujęciu stratygraficznym,  
785 geologicznym i tafonomicznym [The Pleistocene mammal assemblages from the  
786 Geological Museum of PGI-NRI; a stratigraphical, geological and taphonomic approach].  
787 *Przegląd Geologiczny* 65, 53–64.

788

789 **TABLE CAPTIONS:**

790

791 **Table 1.** List of archive sites with remains of the woolly rhinoceros from Poland according to  
792 the order of authors contributions, as of February 2024. NN: locality unknown; NISP: number  
793 of identified specimens; MNI: minimum number of individuals. No data means that the total  
794 data is not yet available.

795 **Table 2.** List of previous results of radiocarbon dating of woolly rhinoceros remains. Data from:  
796 Wojtal (2007), Pawłowska (2012, 2022), Stuart & Lister (2012), Schild (2014), Marciszak et al.  
797 (2019), and modified data from Stefaniak et al. (2023). The symbols ... and – mean that the  
798 date may extend out of range; n/a = not applicable.

799

800 **FIGURE CAPTION:**

801

802 **Fig. 1.** Drawing of a woolly rhinoceros (WOOLRHINOPOLI Project).

803 **Fig. 2.** Mandible (lower specimen) and skull (upper) of a woolly rhinoceros found in 1810 and  
804 1815, respectively in Kamieńczyk site: the first illustration of fossil rhinos from Poland  
805 (modified from Staszic, 1815).

806 Table 1

No.	Site corrected WOOLRHINOPOLI PROJECT	NISP	MNI	Author 1	Author 2	Author 3	Author 4	Author 5	Others
1	Bełchatów	1	no data	Stefaniak et al., 2023 (and some references therein): for each citation here					
2	Bęblowska Dolna Cave, Bęblo	no data	no data	Ossowski 1890	Kowalski 1951, 1959	Geraads et al. 2021	Stefaniak et al. 2023		
3	Bielkowo	2	no data	Kiesow 1880	Schirmacher 1882	Wolff 1903	Kowalski 1959	Geraads et al. 2021	Stefaniak et al. 2023
4	Bielkówko	1	no data	Hermann 1913	Sonntag 1919	Kowalski 1959	Geraads et al. 2021	Stefaniak et al. 2023	
5	Biśnik Cave	79	29	Madeyska 2002	Geraads et al. 2021	Stefaniak et al. 2021, 2023			
6	Bobrów	1	no data	Gürich 1885	Stefaniak et al. 2023				
7	Bogusław	1	no data	Kaczmarek 2004					
8	Bolimów	1	no data	Stefaniak et al. 2023					
9	Borsuka Cave	no data	no data	Wilczyński et al. 2016	Geraads et al. 2021				
10	Bramka Cave	no data	no data	Chmielewski 1975	Geraads et al. 2021				
11	Brodnica	2	no data	Hermann 1913	Sonntag 1919	Kowalski 1959	Geraads et al. 2021	Stefaniak et al. 2023	
12	Brzeziny	no data	no data	Kazanecka 2004	Marciszak et al. 2019	Geraads et al. 2021	Stefaniak et al. 2023		
13	Bychawa	1	no data	Wasilewski 1960	Geraads et al. 2021	Stefaniak et al. 2023			
14	Cave IV, Birów Mountain	no data	no data	Muzolf et al. 2009	Stefaniak et al. 2009	Leshchinskiy 2015	Geraads et al. 2021	Stefaniak et al. 2023	
15	Chmielnik	no data	no data	Król 1998	Geraads et al. 2021	Stefaniak et al. 2023			
16	Chorzów	no data	no data	Römer 1879b	Gürich 1885	Kowalski 1959	Geraads et al. 2021	Stefaniak et al. 2023	
17	Ciemna Cave, Ojców	2	1	Krukowski 1939	Kowalski 1951, 1959	Wojtal 2007	Gradziński et al. 2011	Geraads et al. 2021	Stefaniak et al. 2023
18	Czarkowy	no data	no data	Osmólski 1972	Borsuk-Białynicka 1973	Geraads et al. 2021	Stefaniak et al. 2023		
19	Czarniawka Stream	no data	no data	Stefaniak et al. 2023					
20	Czarnków	1	no data	Lubicz-Niezabitowski 1926	Kowalski 1959	Kaczmarek 2004	Pawłowska 2015b	Geraads et al. 2021	Stefaniak et al. 2023
21	Czechowice-Dziedzice	no data	no data	Konior 1936	Marciszak et al. 2019	Geraads et al. 2021	Stefaniak et al. 2023		
22	Czersk	no data	no data	Stefaniak et al. 2023					
23	Czerwonak near Poznań	2	no data	Lubicz-Niezabitowski 1926	Kowalski 1959	Kaczmarek 2004	Pawłowska 2015b	Geraads et al. 2021	Stefaniak et al. 2023
24	Czerwoniak	no data	no data	Kunisch 1883	Geraads et al. 2021	Stefaniak et al. 2023			
25	Czerwotka	1	no data	Conwentz 1901	Hermann 1913	Kowalski 1959	Geraads et al. 2021	Stefaniak et al. 2023	
26	Dąbrowa Górnicza	no data	no data	Stefaniak et al. 2023					
27	Dąbrówka Mała	no data	no data	Ryzewicz 1933	Stefaniak et al. 2023				
28	Deszczowa Cave	21	4	Cyrek et al. 2000	Sudoł & Cyrek 2015	Wojtal 2007	Geraads et al. 2021	Stefaniak et al. 2023	
29	Dębiec, Poznań	12	no data	Lubicz-Niezabitowski 1926	Rakowski 1933	Wyrwicka 1946	Wasilewski 1960	Kowalski 1959	Kaczmarek 2004 Pawłowska 2015b; Geraads et al. 2021; Stefaniak et al. 2023
30	Długi Most	no data	no data	Hermann 1913	Sonntag 1919	Kowalski 1959	Geraads et al. 2021	Stefaniak et al. 2023	
31	Dobromierz	5	no data	Kaczmarek 2004					

32	Dobrzylin	no data	no data	Ślósarski 1884	Kowalski 1959	Karaszewski 1980	Geraads et al. 2021	Stefaniak et al. 2023	
33	Dzierżno	no data	no data	Heinevetter 1933, 1937	Geraads et al. 2021				
34	Garwolin	4	no data	Żarski et al. 2014	Geraads et al. 2021	Stefaniak et al. 2023			
35	Glinianki Szczęśliwickie, Warsaw	no data	no data	Woroncowa- Marcinowska et al. 2013, 2017	Stefaniak et al. 2023				
36	Główna, Poznań	no data	no data	Lubicz- Niezabitowski 1926	Kowalski 1959	Pawłowska 2015b	Geraads et al. 2021	Stefaniak et al. 2023	
37	Gniew	1	no data	von Baer 1823	Hensche 1860	Müller 1863	Schirmacher 1882	Braun 1910	Hermann 1913; Sonntag, 1919; Kotański 1956; Kowalski 1959; Geraads et al. 2021; Stefaniak et al. 2023
38	Gniewięcin	1	no data	Król 1998	Geraads et al. 2021	Stefaniak et al. 2023			
39	Gołaszyn near Oborniki	1	no data	Lubicz- Niezabitowski 1926	Kowalski 1959	Kaczmarek 2004	Pawłowska 2015b	Geraads et al. 2021	Stefaniak et al. 2023
40	Gorzuchów-Święcko	3	no data	Kunisch 1883	Marciszak et al. 2019	Stefaniak et al. 2023			
41	Gostków	no data	no data	Kunisch 1883	Dathe 1899	Geraads et al. 2021			
42	Góra Kalwaria	2	no data	Stefaniak et al. 2023					
43	Góra Puławska	no data	no data	Krzysztafowicz 1896	Czarnowski 1911b	Kozłowski 1922, 1924	Krukowski 1939	Kowalski 1959	Borsuk-Białynicka 1973; Woroncowa- Marcinowska et al. 2013, 2017; Geraads et al. 2021; Stefaniak et al. 2023
44	Góra Winnica near Kamień Mściowski	no data	no data	Kulczycki 1955	Karaszewski 1976	Woroncowa- Marcinowska et al. 2013, 2017	Geraads et al. 2021	Stefaniak et al. 2023	
45	Grupa	2	no data	Conwentz 1892, 1894	Nehring 1896	Jentzsch 1901	Hermann 1911, 1913	Sonntag 1919	Schroeder 1930; Kowalski 1959; Geraads et al. 2021; Stefaniak et al. 2023
46	Ibramowice	no data	no data	Gürich 1905	Geraads et al. 2021				
47	Izbica	1	no data	Karaszewski 1976	Geraads et al. 2021	Stefaniak et al. 2023			
48	Jankowo near Środa	5	no data	Lubicz- Niezabitowski 1926	Kowalski 1959	Kaczmarek 2004	Pawłowska 2015b	Geraads et al. 2021	Stefaniak et al. 2023
49	Jasna Smoleńska Cave	no data	no data	Sudoł & Cyrek 2015	Sudoł et al. 2016	Geraads et al. 2021	Stefaniak et al. 2023		
50	Jasna Strzegowska Cave	no data	no data	Stuart & Lister 2012	Geraads et al. 2021	Stefaniak et al. 2023			
51	Jelenia Góra	1	no data	Gürich 1885	Kowalski 1959	Marciszak et al. 2019	Geraads et al. 2021	Stefaniak et al. 2023	
52	Jerzmanowice	no data	no data	Wojtal 2007					
53	Józwin	no data	no data	Borsuk-Białynicka 1973	Geraads et al. 2021				

54	Kadyny	1	no data	Conwentz 1909	Hermann 1913	Sonntag 1919	Kowalski 1959	Geraads et al. 2021	Stefaniak et al. 2023
55	Kadzielnia, Kielce	2	no data	Lubicz-Niezabitowski 1926	Kowalski 1954, 1958a, 1959	Pawłowska 2015b	Woroncowa-Marcinowska et al. 2013, 2017	Geraads et al. 2021	Stefaniak et al. 2023
56	Kalinowo	no data	no data	von Baer 1823	Kowalski 1959	Geraads et al. 2021	Stefaniak et al. 2023		
57	Kamieńczyk	2	no data	Staszic 1815	Stefaniak et al. 2023				
58	Karbowo	1	no data	Hermann 1913	Sonntag 1919	Kowalski 1959	Geraads et al. 2021	Stefaniak et al. 2023	
59	Karzec near Gostyń	2	no data	Lubicz-Niezabitowski 1926	Kowalski 1959	Kaczmarek 2004	Pawłowska 2015b	Geraads et al. 2021	Stefaniak et al. 2023
60	Kazimierz, Konin	1	no data	Borsuk-Białynicka 1973					
61	Kłodzko	no data	no data	Otto 1837	Hensel 1852, 1853	Kunisch 1883	Gürich 1885	Kowalski 1959	
62	Kobylnica near Poznań	2	no data	Lubicz-Niezabitowski 1926	Kowalski 1959	Kaczmarek 2004	Pawłowska 2015b	Geraads et al. 2021	Stefaniak et al. 2023
63	Komarowa Cave	43	4	Wojtal 2007	Nadachowski et al. 2009	Geraads et al. 2021	Stefaniak et al. 2023		
64	Komondzianka	no data	no data	Ślósarski 1884	Kowalski 1959	Geraads et al. 2021	Stefaniak et al. 2023		
65	Konin	no data	no data	Borsuk-Białynicka 1973		Stefaniak et al. 2023			
66	Kowanowo near Oborniki	1	no data	Lubicz-Niezabitowski 1926	Kowalski 1959	Kaczmarek 2004	Pawłowska 2015b	Stefaniak et al. 2023	Stefaniak et al. 2023
67	Kowanówko near Oborniki	7	no data	Lubicz-Niezabitowski 1926	Kowalski 1959	Pawłowska 2015b	Geraads et al. 2021	Stefaniak et al. 2023	
68	Koziarnia Cave, Saspów	2	2	Römer 1883, 1884	Czarnowski 1911a	Kowalski 1951, 1959	Szulec 2005	Geraads et al. 2021	Stefaniak et al. 2023
69	Krosinko	in progress	in progress	Pawłowska 2009a, 2009b, 2010, 2015a, 2022	Pawłowska et al. 2022	Pawłowska 2015b	Stefaniak et al. 2023		
70	Kroszyce	no data	no data	Wojtal 2007					
71	Krowodrza, Kraków	2	no data	Jura 1837	Kowalski 1959	Geraads et al. 2021	Stefaniak et al. 2023		
72	Lisia Góra	1	no data	Borsuk-Białynicka 1973	Geraads et al. 2021	Stefaniak et al. 2023			
73	Lubin	no data	no data	Anonymous 1908	Kowalski 1959	Marciszak et al. 2019	Geraads et al. 2021	Stefaniak et al. 2023	
74	Luboń near Poznań	1	no data	Lubicz-Niezabitowski 1926	Kowalski 1959	Kaczmarek 2004	Pawłowska 2015b	Stefaniak et al. 2023	
75	Ludwinów, Kraków	1	no data	Kiernik 1911	Kowalski 1959	Geraads et al. 2021	Stefaniak et al. 2023		
76	Łabędy, Gliwice	no data	no data	Heinevetter 1937	Geraads et al. 2021				
77	Ławy near Siedlce	5	no data	Hrynowiecka et al. 2018	Stefaniak et al. 2023				
78	Łęcze	2	no data	Klien 1910	Hermann 1913	Kowalski 1959	Geraads et al. 2021	Stefaniak et al. 2023	
79	Łokietka Cave	no data	no data	Wojtal 2007	Geraads et al. 2021				

80	Łódź	1	no data	Borsuk-Białynicka 1973	Geraads et al. 2021	Stefaniak et al. 2023			
81	Maciejowice	1	no data	Stefaniak et al. 2023					
82	Malbork	1	no data	Stefaniak et al. 2023					
83	Malta, Poznań	4	no data	Maas 1899	Lubicz-Niezabitowski 1926	Kowalski 1959	Kaczmarek 2004	Pawłowska 2015b	Geraads et al. 2021 Stefaniak et al. 2023
84	Mammutowa Cave, Wierzychowie	no data	1	Zawisza 1878, 1882a, 1882b	Kowalski 1951, 1959	Wojtal 2007	Wojtal et al. 2011	Geraads et al. 2021	Stefaniak et al. 2023
85	Maszycka Cave, Maszyce	no data	no data	Ossowski 1884, 1885	Kowalski 1951, 1959	Lasota-Moskalewska 1993	Geraads et al. 2021	Madeyska & Cyrek 2002	Stefaniak et al. 2023
86	Mechowo near Swarzędz	5	no data	Maas 1899	Lubicz-Niezabitowski 1926	Kowalski 1959	Kaczmarek 2004	Pawłowska 2015b	Geraads et al. 2021 Stefaniak et al. 2023
87	Milowice, Sosnowiec	48	6	Ryzewicz 1933	Kowalski 1959	Marciszak et al. 2019	Geraads et al. 2021	Stefaniak et al. 2023	
88	Minięta	3	no data	Conwentz 1895, 1897	Hermann 1911, 1913	Sonntag 1919	Schroeder 1930	Kowalski 1959	Stefaniak et al. 2023
89	Mroczna Cave	no data	no data	Geraads et al. 2021	Stefaniak et al. 2023				
90	Murek Cave, Czulów	no data	no data	Ossowski 1883	Kiernik 1912	Kowalski 1951, 1959	Geraads et al. 2021	Stefaniak et al. 2023	
91	Na Gaiku II Shelter, Mników	no data	no data	Ossowski 1883	Kowalski 1951, 1959	Geraads et al. 2021	Stefaniak et al. 2023		
92	Na Gołąbcu Cave, Piekary	41	3	Ossowski 1880	Kowalski 1951, 1959	Geraads et al. 2021	Stefaniak et al. 2023		
93	Na Wrzosach Północna Cave, Wrzosey	1	no data	Ossowski 1881	Kowalski 1951, 1959	Geraads et al. 2021	Stefaniak et al. 2023		
94	Nad Galoską Cave, Piekary	no data	no data	Ossowski 1881	Kowalski 1951, 1959	Geraads et al. 2021			
95	Nad Matką Boską Cave, Czulów	no data	no data	Ossowski 1883	Kowalski 1951, 1959	Geraads et al. 2021	Stefaniak et al. 2023		
96	Nad Potoczkiem Cave, Czulów	no data	no data	Ossowski 1883	Kowalski 1951, 1959	Geraads et al. 2021	Stefaniak et al. 2023		
97	Nietoperzowa Cave	8	6	Wojtal 2007	Geraads et al. 2021	Stefaniak et al. 2023			
98	NN_1	no data	no data	Lubicz-Niezabitowski 1926	Pawłowska 2015b				
99	NN_2	no data	no data	Lubicz-Niezabitowski 1926					
100	NN_3	no data	no data	Król 1998					
101	NN_4	1	no data	Borsuk-Białynicka 1973					
102	Nowa Dobra	no data	no data	Conwentz 1905	Hermann 1913	Sonntag 1919	Kowalski 1959	Geraads et al. 2021	Stefaniak et al. 2023
103	Nowawieś Chełmińska	2	no data	Hermann 1913	Stefaniak et al. 2023				
104	Nowe Pole	2	no data	Conwentz 1888	Hermann 1913	Sonntag 1919	Kowalski 1959	Geraads et al. 2021	Stefaniak et al. 2023
105	Oblazowa Cave	7	6	Valde-Nowak & Nadachowski 2014	Geraads et al. 2021	Stefaniak et al. 2023			
106	Oborniki	31	no data	Wahnschaffe 1900a, 1900b, 1914	Krause 1925	Lubicz-Niezabitowski 1926	Schroeder 1930	Kowalski 1959	Kaczmarek 2004 Pawłowska 2015b; Stefaniak et al. 2023

107	Odrzywół	1	no data	Karaszewski 1976	Geraads et al. 2021	Stefaniak et al. 2023			
108	Ojrzanów near Mszczonów	2	no data	Stefaniak et al. 2023					
109	Okiennik Cave, Skarżyce	no data	no data	Lubicz-Niezabitowski 1938	Krukowski 1939	Kowalski 1951, 1959	Pawłowska 2015b	Geraads et al. 2021	Stefaniak et al. 2023
110	Opatów	no data	no data	Król 1998					
111	Oporów, Wrocław	no data	no data	Wiszniowska et al. 2003	Marciszak et al. 2019	Geraads et al. 2021	Stefaniak et al. 2023		
112	Ostróda	no data	no data	Brandt 1877	Kowalski 1959	Geraads et al. 2021	Stefaniak et al. 2023		
113	Ostróg, Racibórz	no data	no data	Marciszak et al. 2019	Geraads et al. 2021	Stefaniak et al. 2023			
114	Ostrzeszów	1	no data	Lubicz-Niezabitowski 1926	Kowalski 1959		Pawłowska 2015b	Geraads et al. 2021	Stefaniak et al. 2023
115	Otmuchów	no data	no data	Römer 1870, 1873, 1879b	Gürich 1885	Kowalski 1959	Marciszak et al. 2019	Geraads et al. 2021	Stefaniak et al. 2023
116	Parchatka	1	no data	Karaszewski 1976	Požaryski 1953	Geraads et al. 2021	Stefaniak et al. 2023		
117	Pawłowiczki	no data	no data	Römer 1879b	Gürich 1885	Pax 1921	Zeuner 1932	Kowalski 1959	Marciszak et al. 2019; Geraads et al. 2021; Stefaniak et al. 2023
118	Perspektywiczna Cave	12	no data	Stefaniak et al. 2023					
119	Perzów	1	no data	Gürich 1885	Kowalski 1959	Marciszak et al. 2019	Geraads et al. 2021	Stefaniak et al. 2023	
120	Piaseczno	1	no data	Majewska 2015	Stefaniak et al. 2023				
121	Płock	no data	no data	Ślósarski 1884	Kowalski 1959	Karaszewski 1980	Geraads et al. 2021	Stefaniak et al. 2023	
122	Pod Kochanką Cave, Mników	no data	no data	Ossowski 1883	Kowalski 1951, 1959	Geraads et al. 2021	Stefaniak et al. 2023		
123	Połom Mountain Cave	no data	no data	Geraads et al. 2021					
124	Poznań	no data	no data	Hermann 1913	Sonntag 1919	Kowalski 1959	Stefaniak et al. 2023		
125	Przechówko	1	no data	Conwentz 1888	Hermann 1913	Sonntag 1919	Kowalski 1959	Geraads et al. 2021	Stefaniak et al. 2023
126	Przezińska Cave, Przegonia	no data	no data	Ossowski 1881	Kowalski 1951, 1959	Geraads et al. 2021	Stefaniak et al. 2023		
127	Przemyków	no data	no data	Zejszner 1856	Kowalski 1959	Geraads et al. 2021	Stefaniak et al. 2023		
128	Przemysł-Jarosław	no data	no data	Zawadzki 1840	Temple 1869	Polanskyj 1928a	Kowalski 1959	Geraads et al. 2021	Stefaniak et al. 2023
129	Pszczółki	1	no data	Hermann 1913	Sonntag 1919	Kotański 1956	Kowalski 1959	Geraads et al. 2021	Stefaniak et al. 2023
130	Pyskowice	no data	no data	Borsuk-Białynicka 1973	Woroncowa-Marcinowska et al. 2013, 2017	Geraads et al. 2021	Stefaniak et al. 2023		
131	Pyskowice-Rzeczyce	no data	no data	Zieliński 1958	Stefaniak et al. 2023				
132	Pyzdry	no data	no data	Geraads et al. 2021					
133	Radłów	1	no data	Stefaniak et al. 2023					
134	Radochowska Cave	no data	no data	Kowalski 1954	Geraads et al. 2021				
135	Raj Cave	70	no data	Kowalski 1972	Majewska 2015	Geraads et al. 2021	Stefaniak et al. 2023		
136	Rataje, Poznań	no data	no data	Lubicz-Niezabitowski 1926	Kowalski 1959	Stefaniak et al. 2023			
137	River Proсна	7	no data	Lubicz-Niezabitowski 1926	Kowalski 1959		Pawłowska 2015b	Geraads et al. 2021	Stefaniak et al. 2023
138	River Vistula (near Warsaw)	1	no data	Borsuk-Białynicka 1973	Stefaniak et al. 2023	Kaczmarek 2004			

139	River Wisła	no data	no data	Jakubowski 1971	Borsuk-Białynicka 1973	Geraads et al. 2021			
140	River Wisznia	no data	no data	Polanski 1928b	Kowalski 1959	Geraads et al. 2021	Stefaniak et al. 2023		
141	Rock Shelter in Strzegowa; Zacisza Cave	no data	no data	Stefaniak et al. 2023					
142	Roznieszew	no data	no data	Stefaniak et al. 2023					
143	Rusko	no data	no data	Gürich 1913	Marciszak et al. 2019	Geraads et al. 2021	Stefaniak et al. 2023		
144	Rybaki-Czartoria	no data	no data	Ruprecht 1976	Geraads et al. 2021				
145	Rzeczyce near Pyskowiec	no data	no data	Kulczycki 1955	Kowalski 1959	Borsuk-Białynicka 1973	Marciszak et al. 2019	Geraads et al. 2021	Stefaniak et al. 2023
146	Sandomierz	1	no data	Stefaniak et al. 2023					
147	Sąpowska Zachodnia Cave	no data	no data	Nadachowski 1988	Geraads et al. 2021	Stefaniak et al. 2023			
148	Shelter above Zegar Cave (Shelter No 388)	no data	no data	Stefaniak et al. 2023					
149	Shelter III (Wilcze I), Sokole Mountain	1	no data	Wojtal 2007	Stefaniak et al. 2009	Geraads et al. 2021	Stefaniak et al. 2023		
150	Shelter V, Złoty Potok	no data	no data	Waga 1853	Ślósarski 1884	Lubicz-Niezabitowski 1913, 1926	Kowalski 1951, 1959	Geraads et al. 2021	
151	Siedliszowice	no data	no data	Pawłowska 2012	Stefaniak et al. 2023				
152	Siemonia	no data	no data	Zieliński 1958	Marciszak et al. 2019	Geraads et al. 2021	Stefaniak et al. 2023		
153	Skarszyn	no data	no data	Römer 1881	Gürich 1885	Pax 1921	Kowalski 1959	Geraads et al. 2021	Stefaniak et al. 2021, 2023
154	Skowarcz	4	no data	Conwentz 1899, 1909	Kumm 1903	Wolff 1905	Staudinger 1908	Hermann 1911, 1913	Sonntag 1919; Schroeder 1930; Kotański 1956; Kowalski 1959; Geraads et al. 2021; Stefaniak et al. 2023
155	Słonne	1	no data	Stefaniak et al. 2023					
156	Sobiecin	1	no data	Stefaniak et al. 2023					
157	Sochaczew	no data	no data	Stefaniak et al. 2023					
158	Splawie	no data	no data	Pawłowska 2017b	Stefaniak et al. 2023				
159	Stajnia Cave	no data	no data	Geraads et al. 2021	Stefaniak et al. 2023				
160	Starogard Gdański	no data	no data	Kumm 1916	Kowalski 1959	Geraads et al. 2021	Stefaniak et al. 2023		
161	Starołęka, Poznań	no data	no data	Maas 1899	Lubicz-Niezabitowski 1926	Kowalski 1959	Pawłowska 2015b	Geraads et al. 2021	Stefaniak et al. 2023
162	Stęszew	2	no data	Kaczmarek 2004					
163	Stradów	1	no data	Stefaniak et al. 2023					
164	Strzegom	no data	no data	Volz 1897	Marciszak et al. 2019	Geraads et al. 2021			
165	Suchanino 1, Gdańsk	no data	no data	Kiesow 1880	Kowalski 1959	Geraads et al. 2021	Stefaniak et al. 2023		



166	Suchanino 2, Gdańsk	1	no data	Schirmacher 1882	Zeise 1903	Hermann 1913	Sonntag 1919	Kowalski 1959	Geraads et al. 2021; Stefaniak et al. 2023
167	Syrzunia	1	no data	Wilke 2004	Marciszak et al. 2019	Geraads et al. 2021	Stefaniak et al. 2023		
168	Szczecin	no data	no data	Stuart & Lister 2012	Geraads et al. 2021	Stefaniak et al. 2023			
169	Śrem	1	no data	Lubicz-Niezabitowski 1926	Kowalski 1959	Pawłowska 2015b	Geraads et al. 2021	Stefaniak et al. 2023	
170	Święcko	no data	no data	Kunisch 1883	Gürich 1885	Kowalski 1959	Geraads et al. 2021	Stefaniak et al. 2023	
171	Świt	no data	no data	Maas 1905	Hermann 1913	Sonntag 1919	Geraads et al. 2021		
172	Targowisko	no data	no data	Jach & Wojtal 2021	Stefaniak et al. 2023				
173	Tatarska Góra	2	no data	Krysiak 1938	Prószyński 1952	Kowalski 1959	Geraads et al. 2021	Stefaniak et al. 2023	
174	Terespol Pomorski	1	no data	Hermann 1913	Sonntag 1919	Kowalski 1959	Geraads et al. 2021	Stefaniak et al. 2023	
175	Tolknicko	no data	no data	Conwentz 1899	Kowalski 1959	Geraads et al. 2021	Stefaniak et al. 2023		
176	Trzebnica	no data	no data	Römer 1888	Marciszak et al. 2019	Geraads et al. 2021	Stefaniak et al. 2023		
177	Tunel Wielki Cave	no data	no data	Nadachowski 1988	Geraads et al. 2021	Stefaniak et al. 2023			
178	Turek	1	no data	Geraads et al. 2021	Stefaniak et al. 2023				
179	ul. Hallera, Wrocław	2	no data	Wiszniowska et al. 2005	Badura & Wiśniewski 2008	Wiśniewski et al. 2009	Stefaniak et al. 2023		
180	ul. Michalska, Przasnysz	no data	no data	Karaszewski 1980	Stefaniak et al. 2023				
181	ul. Spadzista, Kraków	1	no data	Wojtal 2007	Geraads et al. 2021	Stefaniak et al. 2023			
182	ul. Szelałowska, Stare Miasto, Poznań	5	no data	Lubicz-Niezabitowski 1926, 1929	Kowalski 1959	Kaczmarek 2004	Pawłowska 2015b	Geraads et al. 2021	Stefaniak et al. 2023
183	vicinity of Chełm	no data	no data	Prószyński 1952	Kowalski 1959	Geraads et al. 2021	Stefaniak et al. 2023		
184	vicinity of Elbląg	no data	no data	Jentzsch 1878	Wahnschaffe 1909	Kumm 1913	Sonntag 1919	Geraads et al. 2021	
185	vicinity of Morąg	no data	no data	Jentzsch 1878	Schirmacher 1882	Kowalski 1959	Geraads et al. 2021	Stefaniak et al. 2023	
186	vicinity of Ojców	1	no data	Römer 1879a	Kowalski 1959	Wojtal et al. 2011	Geraads et al. 2021	Stefaniak et al. 2023	
187	vicinity of Poznań	1	no data	Maas 1900	Wahnschaffe 1909	Kowalski 1959	Pawłowska 2015b	Geraads et al. 2021	
188	W Dziadowej Skale Cave, Skarżyce	2 or more	2 or more	Kowalski 1958b	Chmielewski 1958	Kowalski 1959	Kaczmarek 2004	Wojtal 2007	Geraads et al. 2021
189	Wadowice	no data	no data	Stach 1956	Kowalski 1959	Borsuk-Białynicka 1973	Geraads et al. 2021	Stefaniak et al. 2023	
190	Walawa	1	no data	Dzieduszycki 1834	Bayger et al. 1914	Polanśkyj 1927	Kowalski 1959	Geraads et al. 2021	Stefaniak et al. 2023
191	Waplewo Wielkie	1	no data	Conwentz 1895, 1899	Hermann 1911, 1913	Sonntag 1919	Kowalski 1959	Geraads et al. 2021	Stefaniak et al. 2023
192	Wielkopolska	no data	no data	Lubicz-Niezabitowski 1926	Pawłowska 2015b				
193	Wiercica Cave	1	no data	Woroncowa-Marcinowska et al. 2013, 2017	Geraads et al. 2021	Stefaniak et al. 2023			
194	Wierzchowska Górna Cave, Wierzchowie	no data	no data	Ossowski 1886, 1887a, 1887b	Kowalski 1951, 1959	Geraads et al. 2021			
195	Wierzenica near Mechowo	1	no data	Lubicz-Niezabitowski 1926	Kowalski 1959	Kaczmarek 2004	Pawłowska 2015b	Geraads et al. 2021	Stefaniak et al. 2023
196	Wilczy Młyn, Poznań	2	no data	Maas 1899	Lubicz-Niezabitowski 1926	Kowalski 1959	Kaczmarek 2004	Pawłowska 2015b	Geraads et al. 2021

197	Wilczyce	179	3	Bratlund 2002	Wiśniewski 2008	Nadachowski et al. 2014	Geraads et al. 2021	Stefaniak et al. 2023	
198	Wilda, Poznań	3	no data	Maas 1899	Lubicz-Niezabitowski 1926	Rakowski 1934	Kowalski 1959	Wasilewski 1960	Kaczmarek 2004 Pawłowska 2015b; Geraads et al. 2021; Stefaniak et al. 2023
199	Wilkowice	no data	no data	Stefaniak et al. 2023					
200	Wola Przemyskowska	1	no data	Borsuk-Białynicka 1973	Geraads et al. 2021	Stefaniak et al. 2023			
201	Września	no data	no data	Stefaniak et al. 2023					
202	Wylotne shelter	7	3	Nadachowski et al. 2015	Geraads et al. 2021	Stefaniak et al. 2023			
203	Zadębcze	1	no data	Prószyński 1952	Kowalski 1959	Stefaniak et al. 2023			
204	Zalesie near Jarocin	3	no data	Behr & Tietze 1911	Hermann 1913	Sonntag 1919	Lubicz-Niezabitowski 1926	Kowalski 1959	Pawłowska 2015b; Geraads et al. 2021; Stefaniak et al. 2023
205	Zawalona Cave	no data	no data	Aleksandrowicz et al. 1992	Geraads et al. 2021	Stefaniak et al. 2023			
206	Zegar Cave	no data	no data	Kowalski 1951 (1846)	Stefaniak et al. 2009	Geraads et al. 2021	Stefaniak et al. 2023		
107	Zegrze	no data	no data	Lubicz-Niezabitowski 1926	Kowalski 1959	Karaszewski 1980	Geraads et al. 2021		
208	Zegrze, Poznań	2	no data	Kaczmarek 2004	Geraads et al. 2021				
209	Zgorzelec	no data	no data	Herr 1924	Heinke 1926	Marciszak et al. 2019	Geraads et al. 2021	Stefaniak et al. 2023	
210	Ziębice	2	no data	Gürich 1885, 1893	Kowalski 1959	Marciszak et al. 2019	Geraads et al. 2021	Stefaniak et al. 2023	
211	Zwoleń	190	4	Gautier 2005	Bratlund 2005	Wojtal 2007	Geraads et al. 2021	Stefaniak et al. 2023	
212	Zygmuntówka	no data	no data	Stefaniak et al. 2023					
213	Żabikowo	2	no data	Lubicz-Niezabitowski 1926	Kowalski 1959	Kaczmarek 2004	Pawłowska 2015b	Geraads et al. 2021	Stefaniak et al. 2023
214	Żerków	1	no data	Kaczmarek 2004					
215	Żmigród	no data	no data	Römer 1873, 1879b	Gürich 1885	Kowalski 1959	Marciszak et al. 2019	Geraads et al. 2021	Stefaniak et al. 2023

814

815

816 Table 2

Site no.	Site name	BP	Calibrated from	Calibrated to	Calibrated median	References	Comments
5	Krosinko	46500 ± 2600	...	46003	-	Pawłowska 2022	excluded; date may extend out of range
6	Nida River near Czarkowy	n/a	n/a	n/a	n/a	Stefaniak et al. 2023	excluded; authors given various data for one dating (13810 ± 70; 13810 ± 700BP) (16750 [18741–14844] cal BP; 16765 [17006–16521] cal BP)
20	Radłów	>45000	n/a	n/a	n/a	Stefaniak et al. 2023	excluded; authors given various data for one dating (n/a for cal BP) (47333 [47984–46795] cal BP)
22	Sochaczew	>49000	...	50663	-	Stefaniak et al. 2023	excluded; date may extend out of range
21	Sąspowska Zachodnia Cave	49000 ± 400	...	50511	-	Stefaniak et al. 2023	excluded; date may extend out of range
17	Wadowice	n/a	n/a	n/a	n/a	Stefaniak et al. 2023	excluded; authors given various data for one dating (16500 ± 100 BP; 16650 ± 100 BP) (19923 [20200–19595] cal BP; 20126 [20412–19867] cal BP)
23	Września	47000 ± 4000	...	45456	-	Stefaniak et al. 2023	excluded; date may extend out of range
24	Zawalona Cave	49000 ± 4000	...	46814	-	Stefaniak et al. 2023	excluded; date may extend out of range
1	Konin	n/a	n/a	n/a	n/a	Stefaniak et al. 2023	excluded; authors given various data for one dating (44000 ± 1400 BP; 44000 ± 140 BP) (46771 [50711–44344] cal BP; 46269 [46835–45838] cal BP)
19	Wilczyce	11400 ± 135	13576	13088	13332	Schild 2014	excluded according to the Stefaniak et al. 2023
11	Skarszyn	13444 ± 226	16949	15601	16275	Marciszak et al. 2019	Excluded, The author changed the date to 16,460 ± 90 BP, 20,126–19,583 cal. BP, Poz-82384 in his latest work (Marciszak et al., 2024).
4	Jasna Strzegowska Cave	16140 ± 90	19805	19180	19493	Stuart & Lister 2012	
7	Nietoperzowa Cave	16780 ± 80	20493	20065	20279	Stefaniak et al. 2023	
4	Jasna Strzegowska Cave	17880 ± 100	22027	21406	21717	Stuart & Lister 2012	
3	Deszczowa Cave	20720 ± 150	25344	24370	24857	Stefaniak et al. 2023	
3	Deszczowa Cave	20800 ± 150	25562	24636	25099	Wojtal 2007	
8	Parchatka near Puławy	21300 ± 130	25887	25292	25590	Stefaniak et al. 2023	
18	Zygmuntówka Quarry near Chęciny	23380 ± 200	27844	27273	27559	Stefaniak et al. 2023	
12	Słonne	24420 ± 170	29107	28149	28628	Stefaniak et al. 2023	
3	Deszczowa Cave	24880 ± 250	29850	28675	29263	Stefaniak et al. 2023	
16	Szczecin	28450 ± 250	33448	31828	32638	Stuart & Lister 2012	
10	Sandomierz	30060 ± 360	35297	33931	34614	Stefaniak et al. 2023	
4	Jasna Strzegowska Cave	29950 ± 220	34849	33996	34849	Cyrek et al. 2016	
3	Deszczowa Cave	31000 ± 400	36174	34610	35392	Lorenc 2013	
3	Deszczowa Cave	31400 ± 400	36550	34840	35695	Wojtal 2007	
9	Perspektywiczna Cave	32100 ± 400	37465	35574	36520	Stefaniak et al. 2023	
9	Perspektywiczna Cave	33900 ± 600	40360	37158	38759	Stefaniak et al. 2023	
15	Stradów	38000 ± 1000	43890	41114	42502	Stefaniak et al. 2023	


5	Krosinko	38500 ± 2000	44074	41621	42848	Pawłowska 2022	
13	Sobiecín	39000 ± 1000	44444	41934	43189	Stefaniak et al. 2023	
2	Biśnik Cave	42200 ± 1400	48298	42882	45590	Stefaniak et al. 2023	
9	Perspektywiczna Cave	42000 ± 1500	48535	42715	45625	Stefaniak et al. 2023	
14	Spławie near Pызdry	42000 ± 2000	51485	42427	46956	Stefaniak et al. 2023	date may extend out of range
21	Sąspowska Zachodnia Cave	>45000	47984	46795	47390	Stefaniak et al. 2023	
24	Zawalona Cave	>45000	47984	46795	47390	Stefaniak et al. 2023	
9	Perspektywiczna Cave	44000 ± 2000	54742	43925	49334	Stefaniak et al. 2023	date may extend out of range
5	Krosinko	48400 ± 32	54759	50055	52407	Pawłowska 2022	date may extend out of range

817

818


A A M

819 Fig.1



820

821 Fig. 2



822

823