

---

**Examples of non-archimedean nuclear Fréchet spaces without a Schauder basis**

by Wiesław Śliwa

*Faculty of Mathematics and Computer Science, A. Mickiewicz University, ul. Matejki 48/49,  
60-769 Poznań, Poland  
e-mail: sliwa@amu.edu.pl*

---

Communicated by Prof. T.A. Springer at the meeting of September 25, 2000

**ABSTRACT**

We solve the problem of the existence of a Schauder basis in non-archimedean Fréchet spaces of countable type (stated in [3]). Using examples of real nuclear Fréchet spaces without a Schauder basis (of Bessaga [1], Mitiagin [5] and Vogt [10]) we construct examples of non-archimedean nuclear Fréchet spaces without a Schauder basis (even without the bounded approximation property).

**1. INTRODUCTION**

In this paper all linear spaces are over a non-archimedean non-trivially valued field  $\mathbb{K}$  which is complete under the metric induced by the valuation  $|\cdot| : \mathbb{K} \rightarrow [0, \infty)$ . For fundamentals of locally convex Hausdorff spaces (lcs) and normed spaces we refer to [8], [6] and [7]. Schauder bases in locally convex spaces are studied in [2], [3], [4] and [9].

Any infinite-dimensional Banach space of countable type is linearly homeomorphic to the Banach space  $c_0$  of all sequences in  $\mathbb{K}$  converging to zero (with the sup-norm) ([7], Theorem 3.16), so it has a Schauder basis. It is also known that any metrizable lcs of finite type has a Schauder basis ([3], Theorem 3.5). In [9] we proved that any infinite-dimensional metrizable lcs contains an infinite-dimensional closed subspace with a Schauder basis.

In this paper we solve the problem stated in [3], whether any Fréchet space of countable type has a Schauder basis. We show that there exist nuclear Fréchet spaces without a Schauder basis. First, we construct an infinite family of pairwise-nonisomorphic nuclear Fréchet spaces with a strongly finite-dimensional

Schauder decomposition but without a Schauder basis (see Theorem 3 and Corollary 5). Next, we give two examples of nuclear Fréchet spaces with a finite-dimensional Schauder decomposition but without a strongly finite-dimensional Schauder decomposition (see Theorem 7 and Corollary 9). Finally, we present an example of a nuclear Fréchet space with a Schauder decomposition but without a finite-dimensional Schauder decomposition (even without the bounded approximation property) (see Theorem 11). Our examples are non-archimedean modifications of the real nuclear Fréchet spaces without a Schauder basis constructed by Bessaga [1], Mitiagin [5], and Vogt [10].

## 2. PRELIMINARIES

The linear span of a subset  $A$  of a linear space  $E$  is denoted by  $\text{lin}A$ .

The linear space of all continuous linear operators from a lcs  $E$  to itself will be denoted by  $\mathcal{L}(E)$ .

A sequence  $(x_n)$  in a lcs  $E$  is a *Schauder basis* of  $E$  if each  $x \in E$  can be written uniquely as  $x = \sum_{n=1}^{\infty} \alpha_n x_n$  with  $\alpha_n \in \mathbb{K}, n \in \mathbb{N}$ , and the coefficient functionals  $f_n : E \rightarrow \mathbb{K}, x \rightarrow \alpha_n (n \in \mathbb{N})$  are continuous.

A sequence  $(P_n)$  of continuous linear non-zero projections on a lcs  $E$  is a *Schauder decomposition* of  $E$  if  $x = \sum_{n=1}^{\infty} P_n x$  for all  $x \in E$  and  $P_n P_m = 0$  for all  $n \neq m$ .

A Schauder decomposition  $(P_n)$  of a lcs  $E$  is *finite-dimensional* if  $\dim P_n(E) < \infty$  for  $n \in \mathbb{N}$ , and *strongly finite-dimensional* if  $\sup_n \dim P_n(E) < \infty$ . Clearly, any lcs  $E$  with a Schauder basis has a strongly finite-dimensional Schauder decomposition.

A lcs  $E$  has the *bounded approximation property* if there exists a sequence  $(A_n) \subset \mathcal{L}(E)$  with  $\dim A_n(E) < \infty$  for  $n \in \mathbb{N}$  such that  $\lim_n A_n x = x$  for all  $x \in E$ . Of course any lcs  $E$  with a finite-dimensional Schauder decomposition has the bounded approximation property.

By a *seminorm* on a linear space  $E$  we mean a function  $p : E \rightarrow [0, \infty)$  such that  $p(\alpha x) = |\alpha|p(x)$  for all  $\alpha \in \mathbb{K}, x \in E$  and  $p(x + y) \leq \max\{p(x), p(y)\}$  for all  $x, y \in E$ . A seminorm  $p$  on  $E$  is a *norm* if  $\text{Ker } p := \{x \in E : p(x) = 0\} = \{0\}$ .

Two norms  $p, q$  on a linear space  $E$  are *equivalent* if there exist positive numbers  $a, b$  such that  $ap(x) \leq q(x) \leq bp(x)$  for every  $x \in E$ . Every two norms on a finite-dimensional linear space are equivalent.

Every  $n$ -dimensional lcs is linearly homeomorphic to the Banach space  $\mathbb{K}^n$ .

A lcs  $E$  is of *finite type* if for each continuous seminorm  $p$  on  $E$  the quotient space  $E/\text{Ker } p$  is finite-dimensional. A metrizable lcs  $E$  is of *countable type* if it contains a linearly dense countable set.

A *Fréchet space* is a metrizable complete lcs.

Any non-decreasing sequence  $(\|\cdot\|_n)$  of norms on a linear space  $E$  defines a metrizable locally convex linear topology on  $E$ . This metrizable lcs will be denoted by  $(E, (\|\cdot\|_n))$ . The sets  $\{x \in E : \|x\|_n \leq m^{-1}\}, n, m \in \mathbb{N}$  form a base of neighbourhoods of 0 in  $(E, (\|\cdot\|_n))$ . A seminorm  $p$  on  $E$  is continuous iff there exist  $m \in \mathbb{N}$  and  $c > 0$  such that  $p(x) \leq c\|x\|_m$  for all  $x \in E$ .

A subset  $B$  of a lcs  $E$  is *compactoid* if for each neighbourhood  $U$  of  $0$  in  $E$  there exists a finite subset  $A = \{a_1, \dots, a_n\}$  of  $E$  such that  $B \subset U + \text{co } A$ , where  $\text{co } A = \{\sum_{i=1}^n \alpha_i a_i : \alpha_1, \dots, \alpha_n \in \mathbb{K}, |\alpha_1|, \dots, |\alpha_n| \leq 1\}$  is the *absolutely convex hull* of  $A$ .

Let  $E$  and  $F$  be locally convex spaces. The linear map  $T : E \rightarrow F$  is *compact* if there exists a neighbourhood  $U$  of  $0$  in  $E$  such that  $T(U)$  is compactoid in  $F$ .

For any seminorm  $p$  on a lcs  $E$  the map  $\bar{p} : E/\text{Ker } p \rightarrow [0, \infty), x + \text{Ker } p \rightarrow p(x)$  is a norm on  $E/\text{Ker } p$ .

A lcs  $E$  is *nuclear* if for every continuous seminorm  $p$  on  $E$  there exists a continuous seminorm  $q$  on  $E$  with  $q \geq p$  such that the map

$$\varphi_{pq} : (E/\text{Ker } q, \bar{q}) \rightarrow (E/\text{Ker } p, \bar{p}), x + \text{Ker } q \rightarrow x + \text{Ker } p$$

is compact. Any nuclear lcs is of countable type ([8], Corollary 4.14).

### 3. RESULTS

First we construct an infinite family of pairwise-nonisomorphic nuclear Fréchet spaces with a strongly finite-dimensional Schauder decomposition but without a Schauder basis (cf [1], [5]).

Put  $\mathbb{N}_0 = \{r \in \mathbb{N} : r > 1 \text{ and } r\mathbf{1} \neq 0 \text{ in } \mathbb{K}\}$ , where  $\mathbf{1}$  is the unit element of  $\mathbb{K}$ . Clearly, the set  $\mathbb{N}_0$  is infinite. Let  $r \in \mathbb{N}_0$ . Let  $\{e_1, \dots, e_r\}$  be a basis of the linear space  $\mathbb{K}^r$  and let  $e_1^*, \dots, e_r^*$  be the coefficient functionals of this basis. Put  $f_1 = \sum_{i=1}^r e_i, f_1^* = \sum_{i=1}^r e_i^*$  and  $f_j^* = e_{j-1}^* - e_j^*$  for  $2 \leq j \leq r$ . It is easy to see that

$$e_1^* = (r\mathbf{1})^{-1} [f_1^* + \sum_{i=2}^r (\sum_{j=2}^i f_j^*)] \text{ and } e_i^* = (e_1^* - \sum_{j=2}^i f_j^*), 2 \leq i \leq r.$$

Hence  $|e_j^*(x)| \leq |r\mathbf{1}|^{-1} \max_l |f_l^*(x)|$  for all  $x \in \mathbb{K}^r, 1 \leq j \leq r$ .

Let  $n \in \mathbb{N}$ . Consider a finite sequence  $(|\cdot|_{n,j}^r)_{j=1}^{r+1}$  of norms on the space  $\mathbb{K}^r$ :

$$|x|_{n,j}^r = \begin{cases} \max(\{2^{n(j-1)}|e_j^*(x)|\} \cup \{2^{nj}|e_i^*(x)| : i \neq j\}) & \text{if } 1 \leq j \leq r, \\ |r\mathbf{1}|^{-1} \max(\{2^{nr}|f_1^*(x)|\} \cup \{2^{n(r+1)}|f_l^*(x)| : l > 1\}) & \text{if } j = r+1. \end{cases}$$

Clearly,  $|x|_{n,j}^r \leq |x|_{n,j+1}^r$  for all  $x \in \mathbb{K}^r, 1 \leq j \leq r$ .

Let  $\beta \in \mathbb{K}$  with  $|\beta| > 1$ . Set  $d = |\beta|$ . For a linear operator  $T : \mathbb{K}^r \rightarrow \mathbb{K}^r, z^* \in (\mathbb{K}^r)^*$  and  $j \leq r+1$  we put  $\|T\|_{n,j}^r = \sup\{|Tx|_{n,j}^r : x \in \mathbb{K}^r, |x|_{n,j}^r \leq 1\}$  and  $\|z^*\|_{n,j}^r = \sup\{|z^*(x)| : x \in \mathbb{K}^r, |x|_{n,j}^r \leq 1\}$ . Then  $|Tx|_{n,j}^r \leq d\|T\|_{n,j}^r|x|_{n,j}^r$  and  $|z^*(x)| \leq d\|z^*\|_{n,j}^r|x|_{n,j}^r$  for all  $x \in \mathbb{K}^r, j \leq r+1$ .

We will need the following

**Lemma 1.** *Let  $(T_k) \subset \mathcal{L}(\mathbb{K}^r)$ . If  $\max_k \dim T_k(\mathbb{K}^r) < r$  and  $\sum_{k=1}^{\infty} T_k x = x, x \in \mathbb{K}^r$ , then  $\max\{\|T_k\|_{n,j}^r : k \in \mathbb{N}, j \leq r+1\} \geq 2^n d^{-2}$ .*

**Proof.** Let  $k \in \mathbb{N}$ . First we show that

$$(*) \quad |e_1^* T_k e_1| \leq \max\{\max_{i \neq j} |e_i^* T_k e_j|, \max_{l > 1} |f_l^* T_k f_1|\}.$$

Let  $t_{i,j} = e_j^* T_k e_i$  for  $1 \leq i, j \leq r$ . Since  $\dim T_k(\mathbb{K}^r) < r$ , then there exist  $\alpha_1, \dots, \alpha_r \in \mathbb{K}$  and  $1 \leq j_0 \leq r$  such that  $\sum_{j=1}^r \alpha_j T_k e_j = 0$  and  $\max_j |\alpha_j| = |\alpha_{j_0}| > 0$ . Hence  $|t_{j_0, j_0}| = |-\sum_{j \neq j_0} \alpha_j^{-1} \alpha_j t_{j, j_0}| \leq \max_{j \neq j_0} |t_{j, j_0}|$ . This yields (\*), if  $j_0 = 1$ .

If  $j_0 > 1$ , then we have

$$\begin{aligned} |t_{1,1}| &= |t_{j_0, j_0} + \sum_{i \neq j_0} t_{i, j_0} - \sum_{i \neq 1} t_{i,1} + (\sum_i t_{i,1} - \sum_i t_{i, j_0})| = \\ &|t_{j_0, j_0} + \sum_{i \neq j_0} t_{i, j_0} - \sum_{i \neq 1} t_{i,1} + \sum_{l=2}^{j_0} f_l^* T_k f_1| \leq \\ &\max\{\max_{i \neq j} |e_i^* T_k e_j|, \max_{l > 1} |f_l^* T_k f_1|\}. \end{aligned}$$

This proves (\*).

$$\begin{aligned} \text{For } 1 \leq i, j \leq r, i \neq j \text{ we have } |e_i^* T_k e_j| &\leq d^2 \|e_i^*\|_{n,j}^r \|T_k\|_{n,j}^r |e_j|_{n,j}^r \leq \\ &d^2 2^{-nj} \|T_k\|_{n,j}^r 2^{n(j-1)} = 2^{-n} d^2 \|T_k\|_{n,j}^r. \end{aligned}$$

$$\begin{aligned} \text{For } 2 \leq l \leq r \text{ we obtain } |f_l^* T_k f_1| &\leq d^2 \|f_l^*\|_{n,r+1}^r \|T_k\|_{n,r+1}^r |f_1|_{n,r+1}^r \leq \\ &d^2 2^{-n(r+1)} |r| \|T_k\|_{n,r+1}^r 2^{nr} \leq 2^{-n} d^2 \|T_k\|_{n,r+1}^r. \end{aligned}$$

Thus  $|e_1^* T_k e_1| \leq 2^{-n} d^2 \max\{\|T_k\|_{n,j}^r : 1 \leq j \leq r+1\}, k \in \mathbb{N}$ . Hence we obtain

$$\begin{aligned} 1 &= \left| \sum_{k=1}^{\infty} e_1^* T_k e_1 \right| \\ &\leq \max_k |e_1^* T_k e_1| \leq 2^{-n} d^2 \max\{\|T_k\|_{n,j}^r : k \in \mathbb{N}, 1 \leq j \leq r+1\}. \end{aligned}$$

This completes the proof of Lemma 1.  $\square$

Put  $\Psi_r = \{(p_1, \dots, p_r) \in \mathbb{N}^r : p_1 < p_2 < \dots < p_r\}$ . Let  $\sigma_r : \mathbb{N} \rightarrow \Psi_r$  be an infinity-to-one surjection (it means that the inverse image of every singleton is infinite).

Let  $n \in \mathbb{N}$ . Set  $p_0 = 0$  and  $\sigma_r(n) = (p_1, \dots, p_r)$ . Consider the following sequence  $(\|\cdot\|_{n,k}^r)_{k=1}^{\infty}$  of norms on the space  $X_n^r = \mathbb{K}^r$ :

$$\|x\|_{n,k}^r = \begin{cases} |x|_{n, j+1}^r & \text{if } p_j < k \leq p_{j+1}, 0 \leq j \leq r-1, \\ |x|_{n, r+1}^r & \text{if } k > p_r. \end{cases}$$

Clearly,  $\|x\|_{n,k}^r \leq \|x\|_{n, k+1}^r$  for  $x \in X_n^r, k \in \mathbb{N}$ .

Let  $\|x\|_k^r = \sup_n n^k \|x_n\|_{n,k}^r$  for  $x = (x_n) \in \prod_{n=1}^{\infty} X_n^r, k \in \mathbb{N}$  and

$$X^r = \left\{ x \in \prod_{n=1}^{\infty} X_n^r : \|x\|_k^r < \infty \text{ for all } k \in \mathbb{N} \right\}.$$

Clearly,  $(\|\cdot\|_k^r)_{k=1}^{\infty}$  is a non-decreasing sequence of norms on the linear space  $X^r$ .

We have the following.

**Proposition 2.** *The metrizable lcs  $X^r = (X^r, (\|\cdot\|_k^r)_{k=1}^\infty)$  is a nuclear Fréchet space with a strongly finite-dimensional Schauder decomposition.*

**Proof.** First we prove that  $X^r$  is a Fréchet space. Let  $(x^m)$  be a Cauchy sequence in  $X^r$  and  $x^m = (x_n^m)_{n=1}^\infty, m \in \mathbb{N}$ . Then

$$(*) \quad \forall k \in \mathbb{N} \forall \epsilon > 0 \exists M(\epsilon, k) > 0 \forall m, l > M(\epsilon, k) \forall n \in \mathbb{N} : n^k \|x_n^m - x_n^l\|_{n,k}^r \leq \epsilon.$$

Hence, for every  $n \in \mathbb{N}, (x_n^m)_{m=1}^\infty$  is a Cauchy sequence in  $(X_n^r, \|\cdot\|_{n,1}^r)$ . Thus  $\lim_m \|x_n^m - x_n^0\|_{n,1}^r = 0$  for some  $x_n^0 \in X_n^r$ . Then  $\lim_m \|x_n^m - x_n^0\|_{n,k}^r = 0$  for all  $k \in \mathbb{N}$ , since  $\dim X_n^r < \infty$ . By  $(*)$  we obtain

$$(**) \quad \forall k \in \mathbb{N} \forall \epsilon > 0 \forall m > M(\epsilon, k) \forall n \in \mathbb{N} : n^k \|x_n^m - x_n^0\|_{n,k}^r \leq \epsilon.$$

Hence  $\forall k \in \mathbb{N} \forall m > M(1, k) : \|\|x^0\|\|_k^r \leq \max\{\|\|x^m\|\|_k^r, \|\|x^m - x^0\|\|_k^r\} < \infty$ , where  $x^0 = (x_n^0)$ . Thus  $x^0 \in X^r$ . By  $(**)$ , we have  $\lim_m \|\|x^m - x^0\|\|_k^r = 0$  for every  $k \in \mathbb{N}$ , so  $x^m \rightarrow x^0$  in  $X^r$ .

To prove that  $X^r$  is nuclear, it is enough to show that for all  $k \in \mathbb{N}, \epsilon > 0$  there exists a finite subset  $A_k^r(\epsilon)$  of  $X^r$  such that  $B_{k+1}^r(1) \subset B_k^r(\epsilon) + \text{co } A_k^r(\epsilon)$ , where  $B_{k+1}^r(1) = \{x \in X^r : \|\|x\|\|_{k+1}^r \leq 1\}$  and  $B_k^r(\epsilon) = \{x \in X^r : \|\|x\|\|_k^r \leq \epsilon\}$ .

Let  $k \in \mathbb{N}, \epsilon > 0$  and  $m \in \mathbb{N}$  with  $m > \epsilon^{-1}$ . For  $n \in \mathbb{N}$  let  $\{e_{n,1}^r, \dots, e_{n,r}^r\}$  be a basis in  $X_n^r$  with  $\|\sum_{i=1}^r \alpha_i e_{n,i}^r\|_{n,k+1}^r \geq \max_{1 \leq i \leq r} |\alpha_i|$  for all  $\alpha_1, \dots, \alpha_r \in \mathbb{K}$ .

Let  $x = (x_n) \in B_{k+1}^r(1)$  and  $x_n = \sum_{i=1}^r \alpha_{n,i} e_{n,i}^r, n \in \mathbb{N}$ . Since  $\|x_n\|_{n,k+1}^r \leq 1$ , then  $x_n \in \text{co}\{e_{n,1}^r, \dots, e_{n,r}^r\}, n \in \mathbb{N}$ . Put  $f_{n,i}^r = (0, \dots, 0, e_{n,i}^r, 0, \dots) \in X^r$ , where  $e_{n,i}^r$  is on  $n$ -th place and  $1 \leq i \leq r, n \in \mathbb{N}$ .

Then  $A_k^r(\epsilon) := \{f_{n,i}^r : n < m, i \leq r\} \subset X^r$  and  $(x_1, \dots, x_{m-1}, 0, 0, \dots) \in \text{co } A_k^r(\epsilon)$ . Since  $n^k \|x_n\|_{n,k}^r \leq n^{-1} \|\|x\|\|_{k+1}^r$  for  $n \in \mathbb{N}$ , then  $(0, \dots, 0, x_m, x_{m+1}, \dots) \in B_k^r(\epsilon)$ . Thus  $x \in B_k^r(\epsilon) + \text{co } A_k^r(\epsilon)$ . Hence  $B_{k+1}^r(1) \subset B_k^r(\epsilon) + \text{co } A_k^r(\epsilon)$ .

For  $m \in \mathbb{N}$  we put  $P_m : X^r \rightarrow X^r, (x_n) \rightarrow (0, \dots, 0, x_m, 0, \dots)$ , where  $x_m$  is on  $m$ -th place. Clearly,  $P_m, m \in \mathbb{N}$ , are continuous linear projections with  $\dim P_m(X^r) = r$  and  $P_m P_l = 0$  for  $m \neq l$ . Let  $x \in X^r$  and  $k \in \mathbb{N}$ . Since  $n^k \|x_n\|_{n,k}^r \leq n^{-1} \|\|x\|\|_{k+1}^r, n \in \mathbb{N}$ , then  $\lim_n n^k \|x_n\|_{n,k}^r = 0$ . Hence  $\sum_{m=1}^\infty P_m x = x, x \in X^r$ . Thus  $X^r$  has a strongly finite-dimensional Schauder decomposition.  $\square$

Now we prove that  $X^r$  has no Schauder basis.

**Theorem 3.** *For any sequence  $(Q_k) \subset \mathcal{L}(X^r)$  such that  $\sum_{k=1}^\infty Q_k x = x, x \in X^r$ , there exists  $k \in \mathbb{N}$  with  $\dim Q_k(X^r) \geq r$ . In particular, the space  $X^r$  has no Schauder basis.*

**Proof.** Suppose, by contradiction, that  $\dim Q_k(X^r) < r$  for any  $k \in \mathbb{N}$ . Since the sequence  $(Q_k)$  is pointwise bounded, then, by the Banach-Steinhaus theorem ([6], Theorem 3.37), the operators  $Q_k, k \in \mathbb{N}$ , are equicontinuous. Hence for

every continuous seminorm  $p$  on  $X^r$  the seminorm  $q: X^r \rightarrow [0, \infty), x \rightarrow \max_k p(Q_k x)$  is continuous on  $X^r$ . Thus there exist integers  $p_0 = 0 < p_1 < \dots < p_r < p_{r+1}$  and a constant  $C$  such that  $\max_k \|Q_k x\|_{p_i+1}^r \leq C \|x\|_{p_i+1}^r$  for  $0 \leq i \leq r$ .

Let  $n \in \sigma_r^{-1}(\{(p_1, \dots, p_r)\})$ . Let  $J_n: X_n^r \rightarrow X^r$  be the natural embedding and  $P_n: X^r \rightarrow X_n^r$  the natural projection. Then  $\|J_n x\|_k^r = n^k \|x\|_{n,k}^r$  for  $x \in X_n^r$ ,  $k \in \mathbb{N}$ , and  $\|P_n x\|_{n,k}^r \leq n^{-k} \|x\|_k^r$  for  $x \in X^r, k \in \mathbb{N}$ . Put  $T_k = P_n Q_k J_n$  for  $k \in \mathbb{N}$ . Then  $(T_k) \subset \mathcal{L}(X_n^r)$ ,  $\max_k \dim T_k(X_n^r) < r$  and  $\sum_{k=1}^{\infty} T_k x = x$  for all  $x \in X_n^r$ . For  $0 \leq i \leq r, x \in X_n^r$  we have

$$\begin{aligned} \max_k \|T_k x\|_{n,p_i+1}^r &= \max_k \|P_n Q_k J_n x\|_{n,p_i+1}^r \leq \max_k n^{-p_i-1} \|Q_k J_n x\|_{p_i+1}^r \leq \\ &n^{-p_i-1} C \|J_n x\|_{p_i+1}^r = C n^{p_i+1-p_i-1} \|x\|_{n,p_i+1}^r \leq C n^{p_i+1} \|x\|_{n,p_i+1}^r. \end{aligned}$$

Since  $\|x\|_{n,p_i+1}^r = \|x\|_{n,p_i-1}^r = |x|_{n,i+1}^r$  for  $x \in X_n^r, 0 \leq i \leq r$ , then

$$\max_k |T_k x|_{n,j}^r \leq C n^{p_r+1} |x|_{n,j}^r \text{ for all } x \in X_n^r, 1 \leq j \leq r+1.$$

Hence  $\max\{\|T_k\|_{n,j}^r: k \in \mathbb{N}, 1 \leq j \leq r+1\} \leq C n^{p_r+1}$ . Using Lemma 1 we obtain  $C n^{p_r+1} \geq 2^n d^{-2}$ . Thus  $2^n n^{-p_r+1} \leq C d^2$  for every  $n$  in the infinite set  $\sigma_r^{-1}(\{(p_1, \dots, p_r)\})$ . Since  $\lim_n 2^n n^{-p_r+1} = \infty$ , we get a contradiction.  $\square$

**Corollary 4.** *Let  $Y$  be a Fréchet space. For any sequence  $(Q_k) \subset \mathcal{L}(X^r \times Y)$  such that  $\sum_{k=1}^{\infty} Q_k z = z, z \in X^r \times Y$ , there exists  $k \in \mathbb{N}$  with  $\dim Q_k(X^r \times Y) \geq r$ .*

*In particular, the Fréchet space  $X^r \times Y$  has no Schauder basis.*

**Proof.** Let  $P: X^r \times Y \rightarrow X^r, (x, y) \rightarrow x$ , and  $S: X^r \rightarrow X^r \times Y, x \rightarrow (x, 0)$ . Put  $Q_k^0 = P Q_k S$  for  $k \in \mathbb{N}$ . Then  $(Q_k^0) \subset \mathcal{L}(X^r)$ ,  $\sum_{k=1}^{\infty} Q_k^0 x = x$  for all  $x \in X^r$  and  $\dim Q_k^0(X^r) \leq \dim Q_k(X^r \times Y), k \in \mathbb{N}$ . Using Theorem 3, we get the corollary.  $\square$

By Theorem 3 and the proof of Proposition 2 we obtain

**Corollary 5.** *The spaces  $X^r, r \in \mathbb{N}_0$ , are pairwise-nonisomorphic.*

Now we construct a nuclear Fréchet space with a finite-dimensional Schauder decomposition but without a strongly finite-dimensional Schauder decomposition (cf [5]).

Put  $\Psi_0 = \bigcup\{\Psi_r: r \in \mathbb{N}_0\}$ . Let  $\sigma_0: \mathbb{N} \rightarrow \Psi_0$  be an infinity-to-one surjection. Let  $n \in \mathbb{N}$ . Set  $\sigma_0(n) = (p_1, \dots, p_{r(n)})$  and  $p_0 = 0$ . Consider the following sequence  $(\|\cdot\|_{n,k}^{r(n)})_{k=1}^{\infty}$  of norms on the space  $X_n^{r(n)} = \mathbb{K}^{r(n)}$ :

$$\|x\|_{n,k}^{r(n)} = \begin{cases} |x|_{n,j+1}^{r(n)} & \text{if } p_j < k \leq p_{j+1}, 0 \leq j \leq r(n) - 1, \\ |x|_{n,r(n)+1}^{r(n)} & \text{if } k > p_{r(n)}. \end{cases}$$

Clearly  $\|x\|_{n,k}^{r(n)} \leq \|x\|_{n,k+1}^{r(n)}$  for all  $x \in X_n^{r(n)}, k \in \mathbb{N}$ .

Let  $\|x\|_k^0 = \sup_n n^k \|x_n\|_{n,k}^{r(n)}$  for  $x = (x_n) \in \prod_{n=1}^{\infty} X_n^{r(n)}$ ,  $k \in \mathbb{N}$  and

$$X^0 = \left\{ x \in \prod_{n=1}^{\infty} X_n^{r(n)} : \|x\|_k^0 < \infty \text{ for all } k \in \mathbb{N} \right\}.$$

Clearly,  $(\|\cdot\|_k^0)$  is a non-decreasing sequence of norms on the linear space  $X^0$ .

The proof of Proposition 2 with obvious changes gives the following.

**Proposition 6.** *The metrizable lcs  $X^0 = (X^0, (\|\cdot\|_k^0))$  is a nuclear Fréchet space with a finite-dimensional Schauder decomposition.*

Similarly, the proof of Theorem 3 with slight changes shows the following theorem.

**Theorem 7.** *For any sequence  $(Q_k) \subset \mathcal{L}(X^0)$  such that  $\sum_{k=1}^{\infty} Q_k x = x$ ,  $x \in X^0$ , we have  $\sup_k \dim Q_k(X^0) = \infty$ . In particular, the space  $X^0$  has no strongly finite-dimensional Schauder decomposition.*

**Corollary 8.** *Let  $Y$  be a Fréchet space. For any sequence  $(Q_k) \subset \mathcal{L}(X^0 \times Y)$  such that  $\sum_{k=1}^{\infty} Q_k z = z$ ,  $z \in X^0 \times Y$ , we have  $\sup_k \dim Q_k(X^0 \times Y) = \infty$ . In particular, the Fréchet space  $X^0 \times Y$  has no strongly finite-dimensional Schauder decomposition.*

Using Proposition 2 and Corollary 4 we obtain the following.

**Corollary 9.** *The Cartesian product  $\prod_{r \in \mathbb{N}_0} X^r$  is a nuclear Fréchet space with a finite-dimensional Schauder decomposition but without a strongly finite-dimensional Schauder decomposition. This space has no continuous norm, so it is not isomorphic to  $X^0$ .*

Finally, we construct a nuclear Fréchet space with a Schauder decomposition but without the bounded approximation property, in particular, without a finite-dimensional Schauder decomposition (cf [10]).

Let  $\alpha \in \mathbb{K}$  with  $0 < |\alpha| < 1$ . For  $k \in \mathbb{N}$  and  $x = (x_{n,p,q}) \in \mathbb{K}^{\mathbb{N} \times \mathbb{N} \times \mathbb{N}}$  we put

$$\|x\|_k = \max \left( \bigcup_{n,p} (\{ |x_{n,p,q}| k^{n+p+q} : q \leq k \} \cup \{ |\alpha^p x_{n,p,q} - x_{n+1,p,q}| k^{n+p+q} : q > k \}) \right)$$

and let

$$X = \{ x \in \mathbb{K}^{\mathbb{N} \times \mathbb{N} \times \mathbb{N}} : \|x\|_k < \infty \text{ for all } k \in \mathbb{N} \}.$$

Note that  $(\|\cdot\|_k)$  is a non-decreasing sequence of norms on the linear space  $X$ . Indeed, if  $x \in X$ ,  $k \in \mathbb{N}$  and  $\|x\|_k = 0$ , then  $x_{n,p,q} = 0$  if  $q \leq k$  and  $x_{n,p,q} = x_{1,p,q} \alpha^{p(n-1)}$  if  $q > k$ . Fix  $p, q \in \mathbb{N}$  and take  $l \in \mathbb{N}$  with  $l > \max\{|\alpha|^{-p}, q\}$ . Then  $|x_{1,p,q}| \max_n (|\alpha^p l|^n) l^{p+q} \leq \|x\|_l < \infty$ . Hence  $x_{1,p,q} = 0$ , so  $x = 0$ .

Since

$\max_{n,p} |\alpha^p x_{n,p,k+1} - x_{n+1,p,k+1}| k^{n+p+k+1} \leq \max_{n,p} |x_{n,p,k+1}| (k+1)^{n+p+k+1}$ ,  
then  $\|x\|_k \leq \|x\|_{k+1}$  for all  $x \in X, k \in \mathbb{N}$ .

We have the following

**Proposition 10.** *The metrizable lcs  $X = (X, (\|\cdot\|_k))$  is a nuclear Fréchet space with a Schauder decomposition.*

**Proof.** First we prove that  $X$  is a Fréchet space. Let  $(x^m)$  be a Cauchy sequence in  $X$  and  $x^m = (x_{n,p,q}^m)_{n,p,q \in \mathbb{N}}, m \in \mathbb{N}$ . Then

$$\forall k \in \mathbb{N} \forall \epsilon > 0 \exists M(\epsilon, k) > 0 \forall m, l > M(\epsilon, k) : \|x^m - x^l\|_k \leq \epsilon.$$

Fix  $n, p, q \in \mathbb{N}$  and  $k \geq q$ . Then

$$\forall \epsilon > 0 \forall m, l > M(\epsilon, k) : |x_{n,p,q}^m - x_{n,p,q}^l| \leq \|x^m - x^l\|_k \leq \epsilon,$$

so  $(x_{n,p,q}^m)_{m=1}^\infty$  is a Cauchy sequence in  $(\mathbb{K}, |\cdot|)$ . Thus  $\lim_m |x_{n,p,q}^m - x_{n,p,q}^0| = 0$  for some  $x_{n,p,q}^0 \in \mathbb{K}$ . Put  $x^0 = (x_{n,p,q}^0)$ . Let  $k \in \mathbb{N}$ . Then

$$\forall \epsilon > 0 \forall m, l > M(\epsilon, k) \forall n, p \in \mathbb{N} \forall q \leq k : |x_{n,p,q}^m - x_{n,p,q}^l| k^{n+p+q} \leq \epsilon.$$

Hence  $\forall \epsilon > 0 \forall m > M(\epsilon, k) \forall n, p \in \mathbb{N} \forall q \leq k : |x_{n,p,q}^m - x_{n,p,q}^0| k^{n+p+q} \leq \epsilon$ .

Similarly, we obtain that  $\forall \epsilon > 0 \forall m > M(\epsilon, k) \forall n, p \in \mathbb{N} \forall q > k :$

$$|\alpha^p (x_{n,p,q}^m - x_{n,p,q}^0) - (x_{n+1,p,q}^m - x_{n+1,p,q}^0)| k^{n+p+q} \leq \epsilon.$$

Hence  $\forall \epsilon > 0 \forall m > M(\epsilon, k) : \|x^m - x^0\|_k \leq \epsilon$ . Since  $\|x^m\|_k < \infty$ , then  $\|x^0\|_k < \infty$ . Thus  $x^0 \in X$ , and  $\lim_m \|x^m - x^0\|_k = 0$  for all  $k \in \mathbb{N}$ .

To prove that  $X$  is nuclear we show that for all  $k \in \mathbb{N}, \epsilon > 0$  there exists a finite subset  $A_k^\epsilon$  of  $X$  such that  $B_{k+1}^\epsilon \subset B_k^\epsilon + \text{co } A_k^\epsilon$  where  $B_{k+1}^\epsilon = \{x \in X : \|x\|_{k+1} \leq 1\}$  and  $B_k^\epsilon = \{x \in X : \|x\|_k \leq \epsilon\}$ .

Let  $k \in \mathbb{N}, \epsilon > 0$ . Choose  $m \in \mathbb{N}$  with  $[k(k+1)^{-1}]^m < \epsilon$ . Put  $t = [k(k+1)^{-1}]$ .

Let  $\{e_{n,p,q} : n, p, q \in \mathbb{N}\}$  be the canonical ‘basis’ in  $X$ . Let  $x \in B_{k+1}^\epsilon$ . For  $(p, q) \in \mathbb{N}^2$  we put  $x^{(p,q)} = \sum_{n=1}^\infty x_{n,p,q} e_{n,p,q}$ ; clearly  $x^{(p,q)} \in B_{k+1}^\epsilon$ .

Let  $p, q \in \mathbb{N}$  with  $p+q < m, q \leq k+1$ . Then  $\|x^{(p,q)} - \sum_{n=1}^{m-1} x_{n,p,q} e_{n,p,q}\|_k \leq$

$$\max_{n \geq m} |x_{n,p,q}| k^{n+p+q} =$$

$$\max_{n \geq m} |x_{n,p,q}| (k+1)^{n+p+q} t^{n+p+q} \leq \|x^{(p,q)}\|_{k+1} t^m \leq \epsilon,$$

and  $\max_n |x_{n,p,q}| \leq \max_n |x_{n,p,q}| (k+1)^{n+p+q} = \|x^{(p,q)}\|_{k+1} \leq 1$ .

Hence  $x^{(p,q)} \in B_k^\epsilon + \text{co } \{e_{n,p,q} : n < m\}$ .

Let  $p, q \in \mathbb{N}$  with  $p+q < m, q > k+1$ . Put

$$y_{n,p,q} = (\alpha^{-np} x_{n,p,q} - \alpha^{-(n+1)p} x_{n+1,p,q}), f_{n,p,q} = \sum_{k=1}^n \alpha^{kp} e_{k,p,q}, n \in \mathbb{N}.$$

Since



$$\sum_{n=1}^{m-1} y_{n,p,q} f_{n,p,q} = \sum_{n=1}^{m-1} e_{n,p,q} (x_{n,p,q} - \alpha^{(n-m)p} x_{m,p,q}),$$

then we obtain  $\|x^{(p,q)} - \sum_{n=1}^{m-1} y_{n,p,q} f_{n,p,q}\|_k =$

$$\begin{aligned} & \left\| \sum_{n=1}^{\infty} x_{n,p,q} e_{n,p,q} + x_{m,p,q} \alpha^{-pm} \left( \sum_{n=1}^{m-1} \alpha^{pn} e_{n,p,q} \right) \right\|_k = \\ & \max_{n \geq m} |\alpha^p x_{n,p,q} - x_{n+1,p,q}| k^{n+p+q} = \\ & \max_{n \geq m} |\alpha^p x_{n,p,q} - x_{n+1,p,q}| (k+1)^{n+p+q} t^{n+p+q} \leq \|x^{(p,q)}\|_{k+1} t^m \leq \epsilon. \end{aligned}$$

Moreover we have  $\max_n \|y_{n,p,q} f_{n,p,q}\|_{k+1} = \|x^{(p,q)}\|_{k+1} \leq 1$ .

Hence  $x^{(p,q)} \in B_k^\epsilon + \text{co} \{\beta_{p,q} f_{n,p,q} : n < m\}$  for any  $\beta_{p,q} \in \mathbb{K}$  with  $\|\beta_{p,q} f_{n,p,q}\|_{k+1} \geq 1$ , for  $n < m$ .

It is easy to check that  $\|x - \sum_{p+q < m} x^{(p,q)}\|_k \leq t^m \|x\|_{k+1} \leq \epsilon$ . Thus  $x \in B_k^\epsilon + \text{co} A_k^\epsilon$  where  $A_k^\epsilon = (\{e_{n,p,q} : n, p+q < m, q \leq k+1\} \cup \{\beta_{p,q} f_{n,p,q} : n, p+q < m, q > k+1\})$ . Hence  $B_{k+1}^1 \subset B_k^\epsilon + \text{co} A_k^\epsilon$ .

For  $n \in \mathbb{N}$  we put  $P_n : X \rightarrow X, x \rightarrow \sum_{p+q=n} x^{(p,q)}$ . Clearly,  $P_n, n \in \mathbb{N}$  are continuous linear projections and  $P_n P_m = 0$  for  $n \neq m$ . Since  $\|x - \sum_{p+q < m} x^{(p,q)}\|_k \leq [k(k+1)^{-1}]^m \|x\|_{k+1}$  for any  $k, m \in \mathbb{N}$ , then  $\sum_{p+q < m} x^{(p,q)} \rightarrow x$  in  $X$ , as  $m \rightarrow \infty$ ; so  $x = \sum_{m=1}^{\infty} P_m x$  for any  $x \in X$ . Of course  $P_n \neq 0, n \in \mathbb{N}$ . Thus  $X$  has a Schauder decomposition.  $\square$

Now we show the following

**Theorem 11.** *The space  $X$  has not the bounded approximation property.*

**Proof.** Suppose, by contradiction, that there exists a sequence  $(A_n) \subset \mathcal{L}(X)$  with  $\dim A_n(X) < \infty, n \in \mathbb{N}$ , such that  $A_n x \rightarrow x$  for all  $x \in X$ . By the Banach-Steinhaus theorem the operators  $A_n, n \in \mathbb{N}$ , are equicontinuous. Thus there exist  $k, l \in \mathbb{N}$  with  $k < l$  and a constant  $C > 0$  such that

$$\max_n \|A_n x\|_1 \leq C \|x\|_k \text{ and } \max_n \|A_n x\|_{k+1} \leq C \|x\|_l \text{ for all } x \in X.$$

Since  $\dim A_n(X) < \infty, n \in \mathbb{N}$ , then

$$\forall n \in \mathbb{N} \exists C_n > 0 \forall x \in X : \|A_n x\|_{k+1} \leq C_n \|A_n x\|_1.$$

Let  $(x^m)$  be a Cauchy sequence in  $(X, \|\cdot\|_l)$  that converges to 0 in  $(X, \|\cdot\|_k)$ . We show that it converges to 0 in  $(X, \|\cdot\|_{k+1})$ . Let  $\delta > 0$ . Then there exist  $t, n, s \in \mathbb{N}$  such that  $\|x^m - x^t\|_l \leq \delta$  for  $m \geq t, \|x^t - A_n x^t\|_{k+1} \leq \delta$ , and  $C_n \|x^m\|_k \leq \delta$  for  $m \geq s$ . Hence, for  $m \geq \max\{t, s\}$  we have

$$\begin{aligned} \|x^m\|_{k+1} & \leq \max\{\|x^m - x^t\|_{k+1}, \|x^t - A_n x^t\|_{k+1}, \|A_n(x^t - x^m)\|_{k+1}, \|A_n x^m\|_{k+1}\} \leq \\ & \max\{\|x^m - x^t\|_l, \delta, C \|x^t - x^m\|_l, C C_n \|x^m\|_k\} \leq \delta(C+1). \end{aligned}$$

Thus  $\|x^m\|_{k+1} \rightarrow 0$ , as  $m \rightarrow \infty$ .

In order to get a contradiction, we take  $p \in \mathbb{N}$  with  $l|\alpha|^p < 1$  and put

$$x^m = \sum_{n=1}^m \alpha^{np} e_{n,p,k+1}, m \in \mathbb{N}.$$

For  $t, m \in \mathbb{N}$  with  $t < m$ , we obtain

$$\|x^m - x^t\|_l = l^{p+k+1} \max_{t+1 \leq n \leq m} (l|\alpha|^p)^n = l^{p+k+1} (l|\alpha|^p)^{t+1}.$$

Thus  $(x^m)$  is a Cauchy sequence in  $(X, \|\cdot\|_l)$ .

Furthermore we have

$$\begin{aligned} \|x^m\|_k &= \max_{1 \leq n \leq m} |\alpha|^p x_{n,p,k+1}^m - x_{n+1,p,k+1}^m |k^{n+p+k+1} = \\ &|\alpha|^{p(m+1)} k^{m+p+k+1} \leq k^{p+k} (l|\alpha|^p)^{m+1} \rightarrow 0, \text{ as } m \rightarrow \infty, \end{aligned}$$

and

$$\|x^m\|_{k+1} = \max_{1 \leq n \leq m} |\alpha|^{pn} (k+1)^{n+p+k+1} \not\rightarrow 0, \text{ as } m \rightarrow \infty. \quad \square$$

**Corollary 12.** *For any Fréchet space  $Y$  the space  $X \times Y$  has not the bounded approximation property.*

#### REFERENCES

1. Bessaga, C. – Nuclear Fréchet Spaces without Bases I. Variation on a Theme of Djakov and Mitiagin. *Bull. Polish Acad. Sci. Math.*, **24**, 471–473 (1976).
2. De Grande-De Kimpe, N. – On the structure of locally  $\mathbb{K}$ -convex spaces with a Schauder basis. *Indag. Mathem.*, **34**, 396–406 (1972).
3. De Grande-De Kimpe, N., J. Kąkol, C. Perez-Garcia and W.H. Schikhof – Orthogonal sequences in non-archimedean locally convex spaces. *Indag. Mathem., N.S.*, **11**, 187–195 (2000).
4. De Grande-De Kimpe, N., J. Kąkol, C. Perez-Garcia and W.H. Schikhof – Orthogonal and Schauder bases in non-archimedean locally convex spaces. *Proceedings of the Sixth International Conference on p-adic Functional Analysis*, Marcel Dekker, New York (to appear).
5. Mitiagin, B.S. – Nuclear Fréchet Spaces without Bases II. The Case of Strongly Finite-dimensional Decompositions. *Bull. Polish Acad. Sci. Math.*, **24**, 475–480 (1976).
6. Prolla, J.B. – Topics in functional analysis over valued division rings. *North-Holland Math. Studies 77*, North-Holland Publ. Co., Amsterdam (1982).
7. Rooij, A.C.M. van – Non-archimedean functional analysis. Marcel Dekker, New York (1978).
8. Schikhof, W.H. – Locally convex spaces over non-spherically complete valued fields. *Bull. Soc. Math. Belgique*, **38**, 187–224 (1986).
9. Śliwa, W. – Every infinite-dimensional non-archimedean Fréchet space has an orthogonal basic sequence. *Indag. Mathem., N.S.*, **11** (3), 463–466 (2000).
10. Vogt, D. – An example of a nuclear Fréchet space without the bounded approximation property. *Math. Z.* **182**, 265–267 (1983).

(Received August 2000)